



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

LANE MEDICAL LIBRARY STANFORD  
D559 .R64 1878  
A manual of anthropometry : or, A guide t  
24503438891

MANUAL  
OF  
ANTHROPOMETRY  
—  
ROBERTS

**LANE**

**MEDICAL**



**LIBRARY**


Gift  
Dr. Horace Gray

12/20/20

12/20/20

61





A

# MANUAL OF ANTHROPOMETRY,

OR

A GUIDE TO THE PHYSICAL EXAMINATION AND  
MEASUREMENT OF THE HUMAN BODY :

CONTAINING A SYSTEMATIC TABLE OF MEASUREMENTS,  
AN ANTHROPOMETRICAL CHART OR REGISTER, AND INSTRUCTIONS FOR  
MAKING MEASUREMENTS ON A UNIFORM PLAN.

*ILLUSTRATED BY NUMEROUS DIAGRAM, CHARTS, AND  
STATISTICAL TABLES.*

LANE LIBRARY

By CHARLES ROBERTS, F.R.C.S.

LATE ASSISTANT SURGEON, VICTORIA HOSPITAL FOR CHILDREN ;  
AUTHOR OF 'THE PHYSICAL DEVELOPMENT AND PROPORTIONS OF THE HUMAN BODY,'  
'THE PHYSICAL REQUIREMENTS OF FACTORY CHILDREN.'



LONDON :

J. & A. CHURCHILL, NEW BURLINGTON STREET.

1878.

[All rights reserved.]

MP



Y&A&B L J&A

LONDON:  
MOBSON AND SONS, PRINTERS, PANCRAE ROAD, N.W.

4551  
R64  
1878

‘THE indispensable part of the experimental observation of physical facts is *the measurement of quantities*. Galileo knew that all physical objects are *extended*, and consequently, by their nature and essence, measurable, though they may not always be measurable by the methods and instruments we possess;—that all physical phenomena take place in periods susceptible of measure—that physical phenomena must be reducible to movements, some perceptible, others imperceptible, by our senses. As regards all these phenomena, he held that the right method was to measure all that was measurable, and to endeavour to render measurable all that was not already directly so. All who have proceeded *à priori*, from Aristotle to Descartes downwards, have arrived at results the falsity of which suffices to condemn their method.’

*Galilée, les Droits de la Science et la Méthode des Sciences Physiques.* Par Th. Henri Martin. (Paris, 1868.)

109265



## PREFACE.

---

THIS small book is a portion of an Introduction to a larger work on the *Physical Development and Proportions of the Human Body*, which I have in preparation; hence its fragmentary character. It is published in this form in order, if possible, to incite a wider interest in an important and interesting field of study hitherto much neglected, and to invite the coöperation of those observers who have already devoted some attention to the subject, with a view to establishing a uniform method of making and recording measurements on the external form of man for scientific purposes.

The Systematic Table of Measurements and the Anthropometrical Chart or Register have been constructed after long and careful consideration of the subject, on anatomical, physiological, and practical grounds; and I trust their fitness for the purposes indicated will be generally admitted. Although at first sight the scheme may seem too elaborate for

general use, I think it will be found on examination, that it contains neither more nor less details than are necessary for the proper study of the subject; it admits, moreover, of being used in part or in its entirety, according to the object the student may have in view, and may, therefore, be accepted as a basis to secure uniformity in making and recording observations, and enabling us to compare together, or combine for statistical purposes, the measurements of different observers.

There are other reasons why I am induced to publish the Table of Measurements and the Chart, which were originally drawn up for my own use. I have been engaged for several years in collecting statistics of measurements of various parts of the body of both sexes, and at all ages, for the purpose of investigating the physical development and proportions of the human body. In the course of my inquiries a large number of the registers of gymnasiums, schools, and public institutions, both military and civil, have passed through my hands, and I have frequently had to regret the imperfection of the details which they contained. This imperfection appears to be due, not so much to carelessness on the part of the keeper of the register or indifference

or unwillingness on that of the pupils to submit to examination, but to ignorance of the kind of facts most deserving of record; to the mode of making the examinations; to the absence of the necessary apparatus; and to the want of a convenient form of book in which to enter the measurements when made. I have had occasion, moreover, to enlist the coöperation of numerous friends in my work, and to write out for each of them precise instructions as to the mode of procedure; hence a series of definitions and directions have been laid down, which I have embodied in the following pages.

To the intelligent traveller, and to many officers of the Army and Navy on foreign stations, this *Manual* and Chart will be found useful, and while supplying a new source of study and occupation, will enable them to contribute to the progress of anthropological science. It has been my good fortune to visit many parts of the world, and to see the various branches of the human race in their own lands; and I have constantly felt the want of some convenient form of register in which to enter the physical proportions of the people; and this want was constantly before my mind while the construction of the Chart was under consideration.

The work will, I trust, be found a useful handbook for many persons who are desirous of assisting the Committee of the British Association for the Advancement of Science, engaged at the present time in investigating the physical proportions and the Racial elements of our British population.

The Systematic Table of Measurements and the Anthropometrical Chart have been adapted to each other; but the Chart is so complete and comprehensive in itself that, after a little practice, reference to the table of measurements will be found unnecessary. It is proposed, therefore, to publish the Chart in separate sheets, for private use or for clinical purposes, or bound in volumes as registers for gymnasiums, hospitals, schools, &c. Exceedingly interesting results would accrue to the individual and to Science if parents would adopt a Chart for each of their children, and record from year to year on their birthdays some of the principal proportions of the body. For this purpose one Chart would suffice for eight or sixteen years, or even for a whole lifetime.

Bolton Row, Piccadilly, W., London,

June 1878.

## WORKS OF REFERENCE ON ANTHROPOMETRY.

---

NOTE.—The following list of works of reference, to which I have made many important additions, is taken from Dr. J. H. Baxter's introduction to *Statistics, Medical and Anthropological, of the Provost-Marshal-General's Bureau* of the United States Government. I am indebted to the courtesy of Dr. Baxter for a copy of this interesting Report, which consists of two thick quarto volumes, and contains the records of the medical examinations and measurements of over a million recruits, &c. during the late civil war in the United States of America.

---

ADAMS (A. L.). On the Physical Requirements of Soldiers. *Brit. and For. Med.-Chir. Rev.* 1875.

ÆLIAN. *Variæ historiæ*. Lib. xiv. 8-16.

AITKEN (DR. WILLIAM). On the Growth of the Recruit and the young Soldier. 12mo. London, 1862.

—— Science and Practice of Medicine. 2 vols. 8vo. London, 1872.

ALBERTI (L. B.). Della architettura di L. Alberti, lib. x.; della pittura, libri iii.; e della statua, libro i.; tradotti in lingua Italiana da Cosimo Bartoli. 3 vols. folio. Londra, 1726.



- ALLEN (NATHAN). Physical Culture in Amherst College. 8vo. Lowell, 1869. [Art. in Appendix, on 'Vital Statistics.']
- ANGERSTEIN. Deutsche Turnzeitung, p. 326. 1864.
- ANTELME. Note sur la céphalométrie. *Mém. de la Soc. d'Anthrop.* t. i. pp. 337-347. Paris, 1860-63.
- ARFE Y VILLAFANE (JUAN). Varia commensuracion para la escultura y arquitectura. Folio. Seville, 1589.
- ARMENINI (J. B.). Veri precetti della pittura, libri iii. 4to. Venice, 1678.
- ARPHE. See ARFE.
- AUDRAN (GERARD). Les proportions du corp humain mesurées sur les belles figures de l'antiquité. Folio. Paris, 1683.
- BARCA (P. A.). Avvertimenti e regole sopra l'architettura civile e militare, la pittura, scoltura e prospettiva. Folio. Milan, 1620.
- BARDON (M. F. DANDRÉ). Traité de peinture. 2 vols. 12mo. Paris, 1765.
- BAXTER (DR. A. M.). Statistics, Medical and Anthropological; Report of the Provost-Marshal-General's Bureau, U.S. Government. 2 vols. 4to. Washington, 1875.
- BEDDOE (DR. JOHN). On the Stature and Bulk of Man in the British Isles. *Mem. Anthrop. Soc.* vol. iii. p. 545. London, 1869.
- On the Physical Characteristics of the Jews. *Trans. Ethnol. Soc.* vol. i. new series, p. 222. London, 1861.
- On the supposed increasing Prevalence of Dark Hair in England. *Anthrop. Review*, vol. i. p. 310. London, 1863.
- On the Head-forms of the West of England. *Mem. Anthrop. Soc.* vol. ii. London, 1865.
- BERGMÜLLER (J. G.). Geometrischer maasstab der säulenordnung und anthropometria, oder natur des menschen. Folio. Augsburg, 1723.
- BERTILLON (A.). De la méthode dans l'anthropologie. *Bull. de la Soc. d'Anthrop.* t. iv. p. 223. Paris, 1863.

- BISCHOFF (T. L. W.). Ueber die veröffentlichten resultate des recrutirungs-geschäftes. 8vo. München, 1867.
- BONOMI (JOSEPH). The Proportions of the Human Figure, according to the ancient Greek Canon of Vitruvius ; also a Canon of the Proportions of the Human Figure, founded upon a Diagram invented by John Gibson, Esq., R.A. 8vo. London, 1857.
- BOSIO (ANT.). Roma sotterranea. Augmentée par Bottari. 3 vols. folio. Rome, 1737.
- BOUCHARDON. Anatomie nécessaire à l'art du dessin. (Publiée par Huguët.) Folio. Paris, 1741. Republished 1802.
- BOUDIN (J. C. M.). Résumé des dispositions légales et réglementaires qui président aux opérations médicales du recrutement, de la réforme et de la retraite dans l'armée de terre. 8vo. Paris, 1854.
- Traité de géographie et de statistique médicales et des maladies endémiques. 2 vols. 8vo. Paris, 1857.
- Etudes ethnologiques sur la taille et le poids de l'homme chez divers peuples. *Recueil de Mémoires de Méd., de Chirurg., &c.* vol. ix. p. 169, vol. x. p. 1. Third series. Paris, 1863.
- BOURGERY (J. M.), BERNARD (C.), et JACOB (N. H.). Traité complet de l'anatomie de l'homme. 8 vols. folio. Paris, 1866-67.
- BOWDITCH (DR. H. P.). The Growth of Children ; Eighth Annual Report of the State Board of Health of Massachusetts. Boston, U.S. 1877.
- BOYD (DR. ROBERT). Tables of the Weight of the Human Body and Internal Organs in the Sane and Insane of both Sexes at Various Ages, arranged from 2614 Post-mortem Examinations. *Phil. Trans.* 1861, pp. 241-262.
- BRENT (W. B.). On the Stature and Relative Proportions of Man at different Epochs and in different Countries. Read before the British Association, Sept. 1844.
- Tables illustrative of the Height, Weight, and Strength of Man. Read before the British Association, June 1845.

- BRENT (W. B.). Comparative Heights of the Soldiers in the British and French Armies, in Proportions of 1000. *The Statistical Companion*, by Banfield & Weld. 16mo. London, 1848.
- BRIGHAM (W. T.). Measurements of 300 Chinese. *Proc. Boston Soc. Nat. Hist.* vol. xi. p. 98. 1866.
- BROCA (PAUL). Recherches sur l'ethnologie de la France. *Mém. Soc. d'Anthrop.* vol. i. pp. 1-56. Paris, 1860.
- Sur l'influence durable de certains croisements de races. *Bull. Soc. d'Anthrop.* vol. i. p. 19. Paris, 1859.
- Sur les proportions relatives du bras, de l'avant-bras et de la clavicule chez les nègres et les Européens. *Ibid.* vol. iii. p. 162. Paris, 1862.
- Sur les projections de la tête, et sur un nouveau procédé de céphalométrie. 8vo. Paris, 1862.
- Nouveaux instruments craniographiques. *Ibid.* 2 sér. vol. iv. p. 101. Paris, 1863.
- Sur le craniographe, &c. *Ibid.* 1 sér. vol. ii. p. 673. Paris, 1861.
- Sur les proportions relatives des membres supérieurs et des membres inférieurs chez les nègres et les Européens. *Ibid.* 2 sér. vol. ii. p. 641. Paris, 1867.
- Nouveau goniomètre. *Ibid.* 1 sér. vol. v. p. 943. Paris, 1864.
- BUFFON (Comte de). Œuvres complètes. 29 vols. 8vo. Paris, 1829-32. (*Histoire de l'homme*, vols. xi.-xiii.)
- BUSK (GEORGE). Observations on a Systematic Mode of Craniometry. *Trans. Ethnol. Soc.* vol. i. new series, p. 341. London, 1861.
- CAMPER (P.). Œuvres de P. Camper qui ont pour objet l'histoire naturelle, la physiologie et l'anatomie. 3 vols. 8vo; and atlas, folio. Paris, 1803.
- CARDAN (J.). Opus novum de proportionibus numerorum, motuum, ponderum, &c. Folio. Bâle, 1570.
- La métoposcopie de Cardan (trad. du Latin) comprise en 13 livres, avec 800 figures de la face humaine, ensemble le

- Traité des signes ou marques naturelles du corps, trad. du Grec de Melampus, par Cl. M. de Laurendière. Folio. Paris, 1658.
- CARUS (C. G.). Die proportionslehre der menschlichen gestalt, zum ersten male morphologisch und physiologisch begründet. Folio. Leipzig, 1854.
- Symbolik der menschlichen gestalt. 8vo. Leipzig, 1858.
- CHAMBERLAINE (J.). Imitations of Original Drawings by Leonard da Vinci, consisting of various drawings of single figures, and, in particular, of very accurate delineations of a variety of anatomical subjects. Folio. London, 1797-1811.
- CHAMPOLLION LE JEUNE (J. F.). Monumens de l'Egypte et de la Nubie. 4 vols. folio. Paris, 1835-45.
- CHAUSSIER (FR.). Recueil anatomique, &c. avec fig. dess. par M. Dutertre. 4to. Paris, 1820.
- COCHIN (C. N.). Œuvres diverses, ou recueil de quelques pièces concernant les arts. 3 vols. 12mo. Paris, 1771.
- CORNEILLE (J. B.). Les premiers éléments de la peinture pratique. 12mo. Paris, 1684.
- COUSIN (JEAN). Livre de pourtraicture par Maistre Jean Cousin, peintre géométrien. 4to. Paris, 1571.
- COWELL. See STANWAY.
- CUMMIN (WILLIAM). Lectures on Forensic Medicine. *London Medical Gazette*, vol. xix. p. 101. 1836.
- DANSON (J. T.). Statistical Observations relative to the Growth of the Human Body (Males) in Height and Weight, from 18 to 30 years of age, as illustrated by the records of the Borough Gaol of Liverpool. *Jour. Stat. Soc.* vol. xxv. p. 20. London, 1862.
- DAVID (EMÉRIC). Recherches sur l'art statuaire chez les anciens et les modernes. 8vo. Paris, 1801-5.
- DAVID (F. A.). Proportions des plus belles figures de l'antiquité, avec figures accompagnées de leur description par Winckelmann. 4to. Paris, 1798.
- DAVIS (DR. B.). *Crania Britannica*. Folio. London, 1856-60.

- DIODORUS SICULUS. *Biblioth. hist. Lib. i. sect. 98.*
- DIONYSIUS OF HALICARNASSUS. *De Isocrates, p. 95.*
- DU GREZ (BERNARD DU PUY). *Traité sur la peinture. 4to. Toulouse, 1699.*
- DURER (ALBERT). *Hierin sind begriffen vier bücher von menschlicher proportion. Folio. Nuremberg, 1528.*
- ELLIOTT (E. B.). *On the Military Statistics of the United States of America. 4to. Berlin, 1863.*
- ELSHOLZ (J. S.). *Anthropometria, sive de mutua membrorum proportione. 4to. Padua, 1654.*
- ELSTER (J. C.). *Die höhere zeichenkunst theoretisch-praktisch, historisch und aesthetisch, &c. 8vo. Leipzig, 1853.*
- FAU (J. A.). *Anatomie des formes extérieures du corps humain à l'usage des peintres et des sculpteurs. 8vo; and atlas, 4to. Paris, 1846.*
- FERGUS (DR. W.), and RODWELL (G. F.). *A Series of Measurements made at Marlborough College. Jour. Anthropological Inst. London, 1874.*
- FIRENZUOLA (AGNOLO). *Dialogo delle bellezze delle donne. 8vo. s.l. 1548.*
- FOCK (H. C. A. L.). *Anatomie canonique, ou le canon de Polyclète retrouvé. 8vo. Utrecht, 1865.*
- *Mémoire sur les proportions du corps de l'homme. Comptes rendus, Acad. des Sci. vol. xxx. p. 661. Paris, 1850.*
- FORBES (DR. JAS. D.). *On the Results of Experiments made on the Weight, Height, and Strength of about 800 Individuals. Reports Brit. Assoc. 1836, part ii. p. 38.*
- GALEN. *De temperamentis, i. 9. De Hippoc. et Platon. decr. iv. 3. De optimi nostri corporis const. i. 2.*
- GALTON (F.). *On the Height and Weight of Boys aged 14 Years in Town and Country Public Schools. Jour. Anthropol. Inst. London, 1875.*
- *Statistics by Intercomparison. Philosophical Magazine. London, 1875.*
- GAURICI (POMPONII). *De sculptura, ubi agitur de symmetriis, de*

- lineamentis, &c. 8vo. Florence, 1504. *See also* GREVIUS. Thesaur. Antiquitat. Græcar. vol. ix. p. 725.
- GAUTIER D'AGOTY (J.). Myologie complète, en 20 pl., avec les explications de Duverney. Folio. Paris, 1746.
- GERDY (P. N.). Anatomie des formes extérieures du corps humain à l'usage des peintres, sculpteurs et dessinateurs. 8vo, plates, folio. Paris, 1830.
- GIBSON (JOHN). *See* BONOMI.
- GILLEBERT D'HERCOURT. Simplification de procédés de mensuration sur l'homme vivant. *Bull. Soc. d'Anthrop.* vol. v. p. 542. Paris, 1864.
- GOULD (B. A.). Investigations in the Military and Anthropological Statistics of American Soldiers. *United States Sanitary Commission Memoirs.* 8vo. New York, 1869.
- GRANVILLE (DR. A. B.). An Essay on Egyptian Mummies. *Philos. Trans.* 1825, p. 269.
- GUIBERT (ADOLPHE). Solution d'une question relative à la probabilité des jugemens rendus à une majorité quelconque. *Liouville Jour. des Mathématiques*, vol. iii. pp. 25-30. 1838.
- Mémoire sur les probabilités des arrêts de deux sortes de cours d'appel. *Comptes rendus, Acad. des Sci.* vol. vii. pp. 650-52. Paris, 1838.
- HAMMOND (DR. W. A.). A Treatise on Hygiene, with special Reference to the Military Service. 8vo. Philadelphia, 1863.
- HAMUSCO. *See* VALVERDE.
- HARGENVILLIERS. Recherches et considérations sur la formation et le recrutement de l'armée en France. 8vo. Paris, 1817.
- HARRISON (J.). Extracts from the Report of the Inspectors of Factories, illustrating the State of Health in the different Factories. *Edin. Med. and Surg. Journal*, 1835.
- HARTING (P.). Le képhalographe. Nouvel instrument destiné à déterminer la figure ou les dimensions du crâne ou de la tête humaine. 4to. Utrecht, 1861.
- HAY (D. R.). The Geometric Beauty of the Human Figure defined:

- to which is prefixed a system of æsthetic proportion applicable to architecture and the other formative arts. 4to. London, 1851.
- HAY (D. R.). *The Natural Principles of Beauty, as developed in the Human Figure.* 8vo. London, 1822.
- On the Science of those Proportions by which the Human Head and Countenance as represented in Works of ancient Greek Art are distinguished from those of ordinary Nature. 4to. Edinburgh, 1849.
- HERSCHEL (SIR J. F. W.). *Essays from the Edinburgh and Quarterly Reviews.* 8vo. London, 1857.
- HUMPHRY (DR. G. M.). *The Human Skeleton.* 8vo. London, 1859.
- HUTCHINSON (JOHN). *Contributions to Vital Statistics, obtained by means of a Pneumatic Apparatus for valuing the Respiratory Powers with relation to Health.* *Quar. Jour. Statist. Soc.* vol. vii. pp. 193-212. London, 1844.
- On the Capacity of the Lungs, and on the Respiratory Functions, &c. *Med.-Chirurg. Trans.* vol. xxix. pp. 137-252. London, 1846.
- Article 'Thorax,' in Todd's *Cyclopædia of Anatomy and Physiology*, vol. iv. p. 1016. London, 1852.
- The Spirometer, the Stethoscope, and the Scale-balance: their Use in discriminating Diseases of the Chest. 8vo. London, 1852.
- IHERING (H.). Ueber das wesen der prognathie und ihr verhältniss zur schädelbasis. *Archiv für Anthropol.* vol. v. p. 359. Brunswick, 1872.
- JACQUART (HENRI). *Mémoire sur la mensuration de l'angle facial, les goniomètres faciaux, et un nouveau goniomètre facial inventé par l'auteur.* *Mémoires lus à la Soc. de la Biologie*, vol. iii. 2 sér. p. 57. Paris, 1856.
- JOHNSTON (A. K.). *The Physical Atlas.* Folio. Edinburgh, 1850.
- JOMARD (E. F.). *Mémoire sur le système métrique des anciens Egyptiens.* Folio. Paris, s.d.

- JOMBERT (C. A.). *Théorie de la figure humaine*, traduit du Latin de Rubens. 4to. Paris, 1773.
- KNOX (ROBERT). *A Manual of Artistic Anatomy*, for the Use of Sculptors, Painters, and Amateurs. 12mo. London, 1852.
- KOPERNICKI (ISIDOR). Ueber den bau der Zigeuner-schädel. Vergleichend-kraniologische untersuchung. *Archiv für Anthropologie*, vol. v. p. 267. 4to. Brunswick, 1872.
- Description d'un nouveau craniographe; étude craniographique des races. *Bull. Soc. d'Anthrop.* 2 sér. vol. ii. p. 559. Paris, 1867.
- KRAUSE (W.). Ueber die aufgaben der wissenschaftlichen kraniometrie. *Achiv für Anthropol.* vol. i. p. 251. Brunswick, 1866.
- LAGNEAU (G.). Notice questionnaire sur l'anthropologie de la France. *Bull. Soc. d'Anthrop.* vol. ii. p. 327. Paris, 1861.
- LAIRESSE (GERARD DE). *Principes du dessin*. Folio. Amsterdam, 1719.
- LAMI (A.). *Anatomie artistique. Myologie superfieielle du corps humain*. Folio. Paris, 1861.
- LAMPREY (—). On a Method of Measuring the Human Form for the Use of Students in Ethnology. *Jour. Ethnol. Soc.* London, 1869.
- LEPSIUS (RICHARD). *Denkmäler aus Aegypten und Aethiopien*. 12 vols. folio. Berlin, 1849-59.
- LICHTENSTEGER (G.). *Arithmétique et géométrie des proportions humaines*. Nürnberg, 1764.
- LIHARZIK (F.). *Das gesetz des wachsthumes und der bau des menschen, die proportionslehre aller menschlichen körpertheile für jedes alter und für beide geschlechter*. Folio. Vienna, 1862.
- *Das quadrat die grundlage aller proportionalität in der natur, und das quadrat aus der zahl 7 die uridee des menschlichen körperbaues*. Vienna, 1865.
- LOMAZZO (G. P.). *Trattato dell' arte, della pittura, scolture ed architettura*. 4to. Milan, 1585.



- LOMAZZO (G. P.). *Traicté de la proportion naturelle et artificielle des choses*, par Jan Pol Lomazzo, peintre Milanois, traduit d'Italien en François par Hilaire Pader, tolosain. Folio. Toulouse, 1649.
- LUCIAN. *De saltatione*, 75; *Philopseudes*, 18; *De morte Peregrini*, 8.
- MACLAREN (A.). *Physical Education and Training*. London, 1866.
- MALLET (EDOUARD). *De la taille moyenne de l'homme dans le canton de Genève. Mémoire lu à la Soc. de Phys. et d'Hist. Nat. de Genève, Déc. 17, 1835.*
- MASCAGNI (PAUL). *Anatomia per uso degli scultori e pittori. Opera postuma*. Folio. Florence, 1816.
- MEDICO (GIUS. DEL). *Anatomia per uso de' pittori ed scultori*. Folio. Roma, 1811.
- MENGES (ANT. RAF.). *Opere in questa edizione corrette ed aumentate de Carlo Fea*. 4to. Roma, 1787.
- MONTABERT (M. P. DE). *Traité complet de la peinture*. 9 vols. 8vo. Atlas, 4to. Paris, 1828-9.
- MORSKOI BORNYK. (Russian Naval Collections.) Vol. xii. 8vo. St. Petersburg, 1871. (Contains results of Physical Examination of Recruits in 1869 and 1870.)
- ORBIGNY (A. D. D'). *Voyage dans l'Amérique méridionale*. 11 vols. 4to. Paris, 1835-47. See vol. iv.
- ORFILA (P.). *Leçons de médecine légale*. 2 vols. 8vo. Paris, 1823. See vol. i. p. 79.
- PAGGI (J. B.). *Definizione et divisione della pittura*. 1607.
- POUSSIN (N.). *Œuvre complet de Poussin, &c.* 8vo. Paris, 1804.
- PLINY. *Histor. nat.* Lib. xxxiv. and xxxv.
- PREISSLER (J. D.). *Die durch theorie erfundene praktik: oder anleitung zu zeichenswerken*. Folio. Nürnberg, 1754.
- PRUNER-BEY. *Questions relatives à l'anthropologie générale. Bull. Soc. d'Anthrop.* vol. v. p. 64. Paris, 1864.
- *Mémoire sur les nègres. Ibid.* vol. i. p. 293. 1859.

- QUETELET (L. A. J.). Recherches sur le poids de l'homme aux différents âges. *Bull. Acad. Sci.* vol. i. p. 20. Bruxelles, 1832-34.
- Sur les Indiens O-Jib-Be-Was, et les proportions de leur corps. *Bull. Acad. Sci.* vol. xiii. p. 70. Bruxelles, 1846.
- Sur les proportions des hommes qui se font remarquer par un excès ou un défaut de taille. *Ibid.* vol. xiv. p. 138. 1847.
- Sur l'étendue superficielle et le volume du corps humain. *Ibid.* vol. xv. part 2, p. 14. 1848.
- Sur les proportions de la race noire. *Ibid.* vol. xxi. p. 96. 1854.
- Sur le constance dans le nombre des mariages, et en la statistique morale en générale. *Ibid.* vol. xxv. p. 89. 1858.
- Sur l'homme et le développement de ses facultés : ou essai de physique sociale. 2 vols. 8vo. Paris, 1835.
- Lettres sur la théorie des probabilités appliquée aux sciences morales et politiques. 8vo. Bruxelles, 1846.
- Anthropométrie ou mesure des différentes facultés de l'homme. 8vo. Bruxelles, 1870.
- QUINTILIAN. *Instit. orator.* Lib. v. 12, xii. 10.
- REGNIER (E.). Mémoire sur le dynamomètre. *Bull. Soc. d'Encourage. pour l'Indust. nat.* vol. xvi. p. 133. Paris.
- REISE der österreichischen fregatte Novara um die erdie, in den jahren 1857, 1858, 1859. 3 vols. 8vo. Vienna, 1861-62.
- REYNOLDS (SIR JOSHUA). *Discourses on Painting.* 4to. London, 1771.
- RÖBER (F. G.). *Elementar-beiträge zur bestimmung der natur-gesetze der gestaltung und des widerstandes, und anwendung dieser beiträge auf natur und alte kunstgestaltung.* 4to. Leipzig, 1861.
- ROBERTS (C.). *The Physical Requirements of Factory Children.* *Journal Statistical Society*, p. 681. London, 1876.
- *The Physical Development and Proportions of the Human Body.* *St. George's Hospital Reports*, p. 1. London, 1874-6.

- ROSELLINI (I.). Monumenti dell' Egitto e della Nubia. 9 vols. 8vo. Plates, 3 vols. folio. Paris, 1832-44.
- RUBENS (P. P.). Théorie de la figure humaine. 4to. Paris, 1773.
- RUSCELLI (GIROLAMO). Le imprese illustri. 4to. Venezia, 1584.
- SABATTINI (G. B.). Tavoli anatomiche per i pittori, scultori ed altri. 4to. Bologna, 1841.
- SALVAGE (J. G.). L'anatomie du gladiateur combattant. Folio. Paris, 1812.
- SANDRART (JOACH. DE). Admiranda sculpturæ veteris. Folio. Norimbergæ, 1630.
- SASSE (A.). Zur wissenschaftlichen kraniometrie. *Archiv für Anthropol.* vol. ii. p. 101. Brunswick, 1867.
- SCHADOW (J. G.). Polyclet oder von den maassen des menschen, nach dem geschlechte und alter mit angabe der wirklichen naturgrösse, &c. Folio and 4to. Berlin, 1834.
- Gottfried Schadow, aufsätze und briefe, nebst einem verzeichniss seiner werke. Zur 100 jährigen feier seiner geburt, 20 Mai 1764, herausgegeben von JUL. FRIEDLAENDER. 8vo. Düsseldorf, 1864.
- SCHERZER (CARL) und SCHWARZ (EDUARD). Ueber körpermessungen als behelf zur diagnostik der menschenracen. *Mittheil. Geogr. Gesellschaft*, vol. iii. p. 11. Vienna, 1859.
- SCHMIDT (CARL). Proportionsschlüssel. Neues system der verhältnisse des menschlichen körpers. 8vo; and plates folio. Stuttgart, 1849.
- SCHWARZ (EDUARD). Anthropology (Novara Expedition). A System of Anthropometrical Investigations as a Means for the Differential Diagnosis of Human Races. 4to. Vienna, 1862.
- SEGOND (L. A.). Procédé de mensuration de la tête applicable à tous les vertébrés, et destiné à découvrir la loi des modifications réciproques entre la face et le crâne, &c. *Compt. rend. Soc. de Biologie*, vol. iii. 2 sér. Paris, 1856.
- SHORTT (JOHN). Notes on Differences in Weight and Stature of Europeans and some Natives of India. *Trans. Ethmol. Soc.* vol. ii. new series, p. 213. London, 1863.

- SIBSON (DR. FRAS.). On the Movements of Respiration in Disease, and on the Use of a Chest-measurer. *Med.-Chirurg. Trans.* vol. xxxi. p. 353. London, 1848.
- SILBERMANN (J. TH.). Proportions physiques ou naturelles du corps humain exprimées en mesures métriques et rapportées à la taille de 1.60 m. *Compt. rend. Acad. des Sci.* vol. xlii. pp. 454, 495 ; vol. xliii. p. 1156. Paris, 1856.
- STANWAY (S.). Results of Investigations made into the Comparative Health and Condition of Factory and Non-factory Children in Manchester and Stockport. Report of Factory Commissioners (Cowell, Chairman). London, 1833.
- STEET (G. C.). Notes on the Development and Growth of Boys between Thirteen and Twenty Years of Age. *St. George's Hospital Reports.* London, 1874-6.
- STORY (W. W.). The Proportions of the Human Figure, according to a new Canon, for Practical Use : with a critical notice of the canon of Polycletus, and of the principal ancient and modern systems. 8vo. London, 1866.
- SUE (J. J.). Sur les proportions du squelette l'homme. *Mém. présentées à l'Acad. Roy. des Sciences*, vol. ii. Paris, 1755.
- TESTELIN (H.). Sentiments des plus habiles peintres sur la pratique de la peinture et de la sculpture. Folio. Paris, 1696.
- THOMSON (A. S.). Observations on the Stature, Bodily Weight; Magnitude of Chest, and Physical Strength of the New Zealand race of men. *Geogr. Soc. Jour.* vol. xxiii. p. 87. London, 1853.
- VALVERDE DI HAMUSCO (GIOV.). Anatomia del corpo humano, &c. (The drawings by Becerra.) Folio. Roma, 1559.
- VAN BREE (MATH.). Leçons du dessin. 8vo. Anvers, 1821.
- VESALIUS (A.). De humani corporis fabrica. Folio. Basil, 1543.
- VILLAFANE. See ARFE.
- VILLERMÉ (L.). Note sur la taille moyenne des habitants de Paris, et sur les proportions des difformités et infirmités qui les

- rendent impropres au service militaire. *Annales des Sciences naturelles*, vol. xi. p. 140. Paris, 1827.
- VILLERMÉ (L.). Mémoire sur la taille de l'homme en France. *Ann. d'Hyg.* vol. i. p. 351. Paris, 1829.
- VINCI (LIONARDO DA). Trattato della pittura. 8vo. Milan, 1804.
- VOGT (CARL). Vorlesungen über den menschen. 8vo. Giessen, 1863.
- VOLPATO (GIOV.). Principi del disegno publicati ed incisi da Giov. Volpato e Rafaele Morghen. Folio. Roma, 1786.
- WATELET (CHR.-HEN.). L'art de peindre. 4to. Paris, 1760.
- WHITE (CHAS.). An Account of the Regular Gradation in Man, and in Different Animals, and from the Former to the Latter. 4to. London, 1799.
- WINCKELMANN (J. J.). Histoire de l'art chez les anciens, traduit de l'allemand. 3 vols. 4to. Paris, 1802.
- WITT (JACOB DE). De proportiën van het menschelijk ligchaam afgebeeld, met de beschrijving, in het Nederduitsch en Fransch. 4to. Amsterdam, s.d.
- WOILLEZ (E. J.). Note sur un nouveau procédé de mensuration de la poitrine. *Archiv. Gén. de la Méd.* 5 sér. t. ix. p. 583.
- WÜLLERSTORF-URBAIR (B. VON, *commodore*). Reise der österreichischen fregatte Novara um die erde in den jahren 1857, 1858, 1859. Anthropologischer theil. 2te abtheilung. Körpermessungen, an individuen verschiedener menschen-racen vorgenommen durch Dr. Karl Scherzer und Dr. Eduard Schwarz, bearbeitet von Dr. A. Weisbach. 4to. Vienna, 1867.
- ZEISING (A.). Neue lehre von den proportionen des menschlichen körpers, &c. 8vo. Leipzig, 1854.
- Die metamorphosen in den verhältnissen der menschlichen gestalt, &c. Folio. Bonn, 1859.

## TABLE OF CONTENTS.



WORKS of reference on Anthropometry . . . . . p. ix.-xxii.

### INTRODUCTION.

Objects and advantages of anthropometric inquiries—Importance to the individual of ascertaining the changes in the form of the body under the influences of age, sex, nurture, occupation, exercise, training, sanitary surrounding, and disease—Social importance of ascertaining the proportions most favourable to health, longevity, and physical endurance, for the insurance of lives and fitness for special duties—Political importance of the subject, as bearing on recruiting for the army and navy, and the employment of children in factories and workshops—Scientific importance of anthropometry: the effect of climate and seasons; the diagnosis of diseases, accidents, and deformities; the classification of the various races of mankind. p. 1-4

### CHAPTER I.

Methods of studying the proportions of the body—Sculptors' and artists' measurements not suitable for scientific purposes—Sir Joshua Reynolds's central form in man—Professor Carus's method—His module equal to a third of the length of the adult spinal column—His statuette of typical proportions—Objections to Professor Carus's *a priori* method of investigation—M. Quetelet's method, based on actual measurements of the living model, and the deduction from them of the mean man (*homme moyen*)—Table and diagram illustrating Quetelet's views—Difference between an average and a mean—Variability of type in different races of mankind—Quetelet's tables based on too small a number of observations, and therefore not reliable . . . . . p. 5-25

## CHAPTER II.

Apparatus—The unit of measurement and of weight and strength—English and metrical systems—Hand-rule, measuring-staff, and measuring-tape—Diagram of scales—Callipers (figure)—Weighing-machine (figure)—Dynamometer (figure)—Figure showing dynamometer arranged as a weighing-machine . . . . *p.* 26-36

## CHAPTER III.

The anthropometrical chart (frontispiece), consisting of a diagram of the body, a register, and a chart for recording observations by the graphic method—Table showing the racial origin of the inhabitants of the British Isles—Classification of occupations—Diseases—Deformities—Classification of the races of mankind . . . *p.* 37-44

## CHAPTER IV.

Systematic table of measurements—Heights—Distances from the vertex (figure)—Diameters and circumferences of head and trunk—Figure showing the movements of the chest by respiration—Measurements of the upper and lower limbs—Weight—Strength . . . *p.* 45-58

## CHAPTER V.

Relative proportions of the body—Relation of the weight to the height—Tables showing average weights for heights—Grouping heights and weights within the same limits—Relation of weight to height shown by interpolation . . . . . *p.* 59-64

## CHAPTER VI.

Construction of statistical tables—Tables showing the height, weight, and chest-girth of the most favoured class; and of artisans living in towns—Height and weight of girls in the United States—Chart-tracings of the heights, breadths, and circumferences of the body—Suitable as scales of proportions for artists . . . *p.* 65-93

## CHAPTER VII.

The physical development and proportions of the body—Influence of race, age, sex, and occupation on the growth of the body—Giants and dwarfs—Proportions of the head; neck; trunk; and the upper and lower extremities—Table of the average height and weight of various classes of the English population . . . . *p.* 94-118

A

## MANUAL OF ANTHROPOMETRY.

---

### INTRODUCTION.

THE objects and advantages of ascertaining the physical proportions of the body, and of the weight and strength of different individuals, are too numerous to receive more than a passing notice here; and I must refer the reader to my published papers, and to numerous foreign works on Anthropometry, for a more complete account of them.\*

On the narrow grounds of individual interest it is desirable to ascertain the changes which take place in the conformation of the body under the various

\* 'The Physical Development and the Proportions of the Human Body,' *St. George's Hospital Reports*, 1874-6; *The Physical Requirements of Factory Children* (London, 1876). See especially Quetelet's *Anthropométrie, ou Mesure des Différentes Facultés de l'Homme* (Bruxelles, 1870); and Schwarz's *Anthropology: a System of Anthropometrical Investigations as a Means for the Differential Diagnosis of Human Races* (Vienna, 1862).



modifying influences of age, nurture, occupation, and sanitary surroundings, and to watch the effects of physical training or athletic exercises — whether favouring health or acting injuriously on the system.

It would be very interesting, on social grounds, to find out, by careful examination, the physical proportions of men who have distinguished themselves by performing great feats of strength, speed, or physical endurance, as such information would indicate which members of a community are best fitted for posts, and could safely undertake duties, requiring the exercise of any of these qualities.

It is important to ascertain what are the physical proportions most favourable to general good health and longevity, or which indicate a disposition to peculiar diseases; whether for sanitary purposes, the insurance of lives, or certifying to the fitness of persons for various occupations at home or abroad.

For the diagnosis of many diseases, accidents, and deformities of the body, physicians and surgeons resort to measurement of the external form, which measurements, to be of value to science, should be made and recorded in a uniform manner.

Many important and interesting physiological and pathological questions bearing on the growth and development of the body can only be set at rest by extensive anthropometrical inquiries, carried out by

a large number of observers in different parts of the world ; such as the effect of climate on the development of a race—our own, for instance, in England, America, Australia, India, &c.—the effects of the season of the year, of peculiar hygienic conditions, and of diseases of hereditary or accidental character ; and it is absolutely necessary that all such observations should be made and recorded in an easily recognised and systematic manner.

From a wider and national point of view it is desirable that we should know the rate of growth of children from year to year, the proportions which exist at different ages, the period at which the body attains maturity, and recognise the nature and peculiar effects of various influences at work in modifying the physical development of our labouring population ; to enable us to regulate the employment of children in factories and workshops, to guide us in the selection of recruits for the military and naval services of the country, and to determine, if possible, the much-debated question of the physical degeneracy of a people.

Naturalists have arrived at the conclusion that measurement of the different dimensions of the body constitutes the best means of classifying the various races and varieties of mankind. ‘Detailed research more and more justifies Blumenbach’s words, that

innumerable varieties of mankind run into one another by insensible degrees. This state of things, due partly to mixture and crossing of races, and partly to independent variations of type, makes the attempt to arrange the whole human species within exact bounded divisions an apparently hopeless task. It does not follow, however, that the attempt to distinguish special races should be given up; for there, at least, exist several definable types, each of which so far prevails in a population as to be taken as its standard. It is by following M. Quetelet's method of defining such types that the subdivisions of mankind into races, so far as it has been done to any purpose, has been carried out by anthropologists.\*

\* Dr. E. R. Tylor, in *Ency. Brit.*, 9th edit., article 'Anthropology.'

## CHAPTER I.

### METHODS OF STUDYING THE PROPORTIONS OF THE HUMAN BODY.

**W**HEN we take a superficial view of the human species we find it composed of individuals varying so widely in physical character that it appears almost hopeless to subject it to a methodical and scientific study. On closer examination, however, certain typical forms are found to prevail through all the stages of development round which the irregularities of size and weight group themselves in the most uniform manner.

The proportions of the body have been studied by sculptors and artists from the earliest times of which we have any record, and, as will be seen from the foregoing list of works of reference, numerous books have been written by them on the subject. Indeed, until quite recently, the study has been appropriated by them, and neglected by physiologists and naturalists as one pertaining only to the Fine Arts. Their works are, however, of little scientific value, as we are ignorant for the most part how the proportions were determined; whether according to

a vague idea of the beautiful in the human form or from actual measurements of the living body. M. Quetelet, who has studied these early records with the eye of an artist as well as a scientific inquirer, thus speaks of them: 'I have tried to gain a glimpse of the principal works on the proportions of man by consulting different times and different peoples. I have been astonished, I avow, at the few original documents I have found. Even according to these, the authors employed but few models to fix the size and the beauty of the conformation of the members, and oftenest the proportions of the child were totally wanting. We see, however, from the works of the Greeks alone, who have always remained masters in these matters, the infinite precautions which they took to arrive at their ends, and to unite elegance of proportion to exactitude of form. Imbued with a knowledge of these, they indeed only consulted the stature of man for the assemblage and agreement of the different parts. Phidias, it is said, to arrive at elegance, employed twenty models; he borrowed from each of them the most beautiful parts, his knowledge of the human form permitting him to arrange them with all the necessary strength and dignity.\*

It is foreign to my object in this small work to describe the various attempts which have been made

\* *Anthropométrie*, p. 412.

to reduce the study of human proportions to exact schemes by the various authors to whom M. Quetelet refers. From the *Silpi Sastri* of the Sanscrit manuscript of India to Mr. Story's disquisition on the mystical qualities of triangles, squares, and circles (1866), the same fallacy pervades them all, namely, a belief that the key to the theory is to be found in the occult relations of numbers or in the parts of a geometrical diagram.\*

There is one artist, however, to whom this remark does not apply, namely, our illustrious countryman Sir Joshua Reynolds, who seems to have had a clear conception in his own mind of the true theory of proportions, although it does not appear that he took any pains to establish it by actual measurements.

As science owes to the poetical mind of Goethe the discovery which has so greatly simplified the study of vegetable morphology—that the leaf is the typical form of the plant, and that all the other organs are modifications of the leaf—so she owes to the æsthetic mind of Sir Joshua Reynolds the idea of the existence of a typical form in man, and the order which prevails in the seeming variations from that type. In this, as in so many other matters, Sir

\* See Dr. Baxter's Introduction to *Statistics, Medical and Anthropological*, p. lxix.

Joshua was in advance of his times, and it remained for M. Quetelet to reduce this artistic conception to a scientific generalisation, and to demonstrate its truth and utility by wide and careful observation on living models.

Sir Joshua's views on this subject form the substance of his third discourse, delivered to the students of the Royal Academy of Arts, in 1770, of which he was then President. Although the language in which they are expressed is such as suited his own tone of mind, and was fitted for an audience of art-students, it is sufficiently exact and interesting to deserve the attention of scientific men. 'All the objects which are exhibited to our view by Nature,' says he, 'upon close examination will be found to have their blemishes and defects. The most beautiful forms have something about them like weakness, minuteness, or imperfection. But it is not every eye that perceives these blemishes. It must be an eye long used to the comparison of these forms; and which, by a long habit of observing what any set of objects of the same kind have in common, has acquired the power of discerning what each wants in particular. By this means we acquire a just idea of beautiful forms; we correct Nature by herself, her imperfect state by her more perfect, and make out an abstract idea of forms more perfect than

any one original. . . . From reiterated experience and a close comparison of the objects of Nature, the artist becomes possessed of a central form from which every deviation is deformity. . . . To the principle I have laid down, that the idea of beauty in each species of being is an invariable one, it may be objected that in every particular species there are various central forms, which are separate and distinct from each other, and yet are undoubtedly beautiful; that in the human figure, for instance, the beauty of Hercules is one, of the Gladiator another, of Apollo another; which makes so many different ideas of beauty. It is true, indeed, that these figures are each perfect in their kind; but still none of them is the representation of an individual, but of a class. And as there is one general form which belongs to the human kind at large, so in each of these classes there is one common idea and central form which is the abstract of the various individual forms belonging to that class. Thus, though the forms of childhood and age differ exceedingly, there is a common form in childhood and a common form in age, which is the more perfect as it is more remote from peculiarities. But I must add further, that though the most perfect forms of each of the general divisions of the human figure are ideal, and superior to any individual form of that



class, yet the highest perfection of the human figure is not to be found in any one of them. It is not in Hercules, nor in the Gladiator, nor in the Apollo ; but in that form which is taken from them all, and which partakes equally of the activity of the Gladiator, of the delicacy of the Apollo, and the muscular strength of the Hercules. . . . There is, likewise, a kind of symmetry or proportion which may properly be said to belong to deformity. A figure lean or corpulent, tall or short, though deviating from the type, may still have a certain union of the various parts which may contribute to make them on the whole not unpleasing.'

Before proceeding to explain how M. Quetelet has established the truth and utility of these views of the artist, and reduced them to a scientific form, I must notice the method of studying and defining the proportions of man adopted by Professor Carus of Dresden.

This distinguished physiologist, considering the proportions of man as an object of morphology, has tried to find physiological laws to fix what he calls the 'canon,' or, to use an expression familiar to architects in the case of the column, the 'module' of our organisation. For this purpose he supposes a figure, the model of which he had executed by the sculptor Rietochel according to his injunctions, and

after admitting the variations of certain parts pertaining to the sexes, he deduces from it the forms both of man and woman.

‘If people are anxious,’ says he, ‘to find the true key of our proportions, they should start from the vertebral column, which is, so to speak, the real organic ell, divided into 24 inches (free vertebræ).

‘When we open the ovum of a mammiferous animal at the commencement of its incubation, we find as the first model of the future animal the germinative disc, intersected in the middle by a line which subsequently becomes the dorsal column or spine. . . . This line becomes longer, and we observe as an already nearly complete model of the future animal a division of this line by the rudiments of the vertebræ. Properly speaking, this form is, then, the first “canon” of every other organisation of the future framework, for according to its method of production and development the whole organisation has to be regulated.

‘Now there are extremely interesting relations when we examine the ratios of length of the free vertebral column in the new-born infant and the adult. In the case of the first (that is to say, at the end of fœtal life), it will be found that the length of all the twenty-four free vertebræ corresponds, in

the normal child, very exactly to one-third of the length of the same free vertebral column, of twenty-four vertebræ, measured in the adult, at the period of the cessation of growth, by a perpendicular line from the *processus spinosus atlantis* to the *processus spinosus lumbaris ultimæ*.' This standard, or, as Professor Carus calls it, 'modulus' of a third of the length of the adult spine, he considers to be formed and established on a physiological and philosophical basis, and its application gives the exact mathematical means of the proportions of our organism, exact in the strictest sense of the word, and therefore ideal proportions which can never be entirely realised in human nature. The following are some of the measurements given by Professor Carus: The length of the skull from the forehead to the occiput, and the height of the vertex from the lower margin of the upper jaw, are each equal to one module; the circumference of the head equals three modules. The length of the sternum, the length of the scapula, and the length of the hand are each equal to one module; the arm and forearm are equal to two modules; the thigh two and a half, the leg two, and the foot from the ankle to the tip of the toe one module. The height of the whole body is nine and a half modules. The module for the adult measures eighteen centimètres, or rather more than seven inches. Dr.

Humphry,\* who has made measurement of numerous *skeletons* for the purpose of testing the value of this means of determining the scale of the proportions of the figure, found them in a general manner confirmatory of the result obtained by Professor Carus; but the exceptions to the rule were very numerous.

Professor Carus gives the following account of his statuette and the mode of using it: 'No sex has been assigned to this little statue; and it is easy to see that, in order to form a living individuality, the modulus or canon must always be made to vary slightly. For instance, if I wished to depict a woman's body, I should give a little less breadth to the shoulders, and I should make some members rather more voluminous; while I should act exactly the contrary in the case of a man. In the same way the individualities might be varied: if I wished to represent a Cicero or a Leibnitz, I should give to the head more than a module in height and length, and less at the extremities; on the other hand, if I wished to represent an athlete or a giant, I should add to the limbs, and should take ten or more modules as the height of the whole body. By this means one could even succeed in depicting every sort of expression by an algebraical formula, where one would have the

\* *On the Human Skeleton.*

same elements, but increased or diminished in their value.'

Thus it appears that Professor Carus uses his 'canon' either as a kind of artist's lay-figure, which he dresses out according to his fancy, or as a skeleton, which he clothes with flesh according to his anatomical and physiological knowledge—knowledge, it must be remembered, which must be first obtained from actual observation and measurement of the living model. The 'canon' may indeed be theoretically correct, but it can be of little practical use for scientific purposes. The greater breadth of shoulders required to convert the statue into the figure of a man must first be determined by actual measurement, as must also the greater breadth of pelvis to convert it into the form of a woman, before we can be satisfied that it represents the natural human form. The difficulty would be still greater if it were attempted to represent any decided variation from the typical form. In the case of a giant, for instance, it is not sufficient to add half a module in equal proportions to the nine and a half modules representing the stature of an ordinary man in order to produce the giant; for actual observation and measurement show that the size of the head and trunk of giants differs little from those of men of ordinary stature, and that the excess of height of the former is chiefly due to

an unusual development of the lower extremities relatively to the rest of the body. Professor Carus's canon, moreover, renders no assistance to the study of the progressive development of the body, as we know that the different parts of the body develop at various rates. Thus, in the young child the middle point is near the navel, but in the adult man it is below the pubes.

M. Quetelet's method of studying the development and proportions of the human body will most strongly recommend itself to the English student, as it is based on the inductive, and not, like Carus's, on the deductive method of investigation, and applies, moreover, to the whole period of the life of man. This distinguished mathematician, like the artist and the physiologist, recognises the existence of a central or typical form in man; but while theirs are ideal forms, his is a real one, and is the *mean* result of a large number of actual measurements of the living body. Hence the 'central form' of Sir J. Reynolds and the 'canon' of typical proportions of Professor Carus become the 'mean man' (*homme moyen*) of M. Quetelet.

'When I first began my studies,' says M. Quetelet,\* 'concerning the proportions of the human body, I was dismayed at the extent which this immense

\* *Anthropométrie*, p. 13.

field of research presented. In fact, I came to the conclusion that in order to obtain data which were fairly complete, I should have to determine for each individual a considerable number of numerical elements; and I was not as yet certain as to how many individuals I should have to measure in order to discover the characteristic details of any particular age. As it was necessary, moreover, to repeat this labour for both sexes, and to follow, so to speak, step by step, the shades of difference which distinguish each period of existence, I came to the conclusion that the life-long labours of the most patient observer would prove inadequate to the task. . . . . Fortunately, I soon perceived that my fears were exaggerated, and that my labour would be greatly simplified by the discovery of a principle which would serve as a basis for it, and which I think I may consider as the most curious and interesting result at which I have arrived.

‘A large number of naturalists and philosophers have attempted to prove, by a course of reasoning which is more or less conclusive, *the unity of the human species*. I believe that I have succeeded in demonstrating, not only that this unity exists, but that our race admits of a type or model the different proportions of which can be easily determined.

‘If there were an absence of type, and if men


were unlike one another, not from the effects of accidental causes, but because no common law really existed among them, they might be measured, as regards height for instance, without all the individual measurements offering any particular character or any definite numerical relation.

‘If, on the contrary, all men have to a certain degree been cast in the same mould, and if they issue from it with differences which are purely accidental, the groups will no longer be formed in an erratic manner; but their numerical values, in accordance with the theory of probabilities, will be subject to preëstablished laws, so that the numbers which represent each can be determined *à priori*. There exist, therefore, for this entirely special case, characteristics by which we may recognise whether individuals belong to the same type, and only differ owing to fortuitous causes. Another consequence of the theory is, that the greater the number of observations the more do fortuitous causes explain each other, and make the general type, which they at first tend to screen, stand out prominently. Thus, in the human race, when individuals only are considered, all heights are met with, at least within certain limits; those who come nearest the average are the most numerous; those who deviate the most from it form the smallest number; and the groups follow numerically



a law (the binomial law of Newton\*) which may be laid down beforehand. In the case of man this law is verified, not only in relation to the entire height, but also as regards his various members; and the same is the case with the weight, or strength, or any quality which can be measured and reduced to numbers.'

The following table and diagram, which represent the heights from actual measurement of 430 English public-school boys of the age from eleven to twelve years, will illustrate Quetelet's views and make them more easily understood:



\* 'This law for the mathematician is nothing but the law of the coefficients of the binomial quantity developed, which, for the sake of brevity, we will call *the binomial law*. If, for instance, *four* balls are taken out of an urn which contains a considerable number

It will be seen that the numbers arrange themselves according to a very uniform rule, the most numerous groups are in the middle of the column at 53 and 54 inches, while the groups at 52 and 55 inches are less in number, and those at 51 and 56 inches are still fewer, and so on till the extremely small number of the very short and very tall boys of 47 and 60 inches is reached. It is thus ascertained that the mean or typical boy of the class and age given is 53.5 inches, and as representing the most numerous group he forms the standard from which the other groups of boys decrease in number as they depart further and further from his proportions.

In the diagram, which has been drawn to a scale, the length of the horizontal lines (*abscissæ*) represents the numbers of boys in each group, and the curved line binding the ends together is the well-

---

of white balls and exactly the same number of black balls, we shall, if we call a white ball  $a$  and a black ball  $b$ , get the following formula as the result of drawing :

$$a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4.$$

The sum of the coefficients of the terms indicates the number of drawings which can take place; *i.e.* a single one produces four white balls, four producing three white balls and one black, six producing two white and two black balls, &c., or in all sixteen possible chances for each drawing. Each of these sixteen drawings given will consequently, according to the laws of probabilities, take place once in sixteen drawings.' *Anthrop.* p. 16.

known 'binomial curve,' or the 'curve of the frequency of error.' Now it will be seen that this curve is not quite uniform, and that the lower half (from the mean to the dwarfs) is less regular than the upper; and if the numbers are counted it will be found that there are ten more boys below the mean than above it, consequently the *average* (obtained in the usual way, by dividing the sum of the values observed by the number of observations) is lower than the *mean*, which is represented by the largest group. The difference in this instance is not very great, the average being 53·43, and the mean 53·5 inches, but in some instances it is much greater; and it is exceedingly important that the difference here indicated should be borne in mind, for in it consists much of the practical value of Quetelet's method.

Dr. Tylor, in the article already referred to, gives a practical illustration of the difference between a mean and an average when employed to separate two races in a mixed population. 'The population of a Californian settlement,' says he, 'made up of whites and Chinese, might show two predominant groups (one of 5 feet 8 inches, the other of 5 feet 4 inches), corresponding to the two racial types. It need hardly be said that this method of determining the mean type of a race, as being that of its really existing and most numerous class, is altogether superior

to the mere calculation of an average, which may actually be represented by comparatively few individuals, and those the exceptional ones. For instance, the average stature of the mixed European and Chinese population just referred to might be 5 feet 6 inches, a worthless and, indeed, misleading result.'

M. Quetelet very properly insists on this difference between an average and a mean, and Sir J. Herschel thus refers to it (*Edin. Rev.* 1850): 'An average may exist of the most different objects; as of the height of houses in a town, or the sizes of books in a library. It may be convenient to convey a general notion of things, but it involves no conception of a natural and recognisable central magnitude, all differences from which ought to be regarded as deviations from a standard. The notion of a mean, on the other hand, does imply such a conception, standing distinguished from an average by this feature, namely, the regular march of the groups, increasing to a maximum, and then again diminishing. An average gives us no assurance that the future will be like the past; a mean may be reckoned on with the most implicit confidence. All the philosophical value of statistical results depends on a due appreciation of its consequences.'

I have dwelt on the importance of recognising

the difference between a mean and an average when applied to the study of the physical proportions of man, because it has been overlooked by writers in this country, and because I propose to use it further on, when treating of the relation between the height and weight of man.

All the published statistics relating to this subject which have come under my notice are simple *averages*, and not *means*. But this must necessarily be the case when a small number of observations is dealt with, as was the case with M. Quetelet, who, after measuring thirty individuals of the same age, divided them into three groups, and finding the groups so nearly alike, concluded that the measurement of ten 'regularly constituted' models would suffice to determine the typical proportions for each age and sex. In this respect my tables show that M. Quetelet was in error, and to this must be attributed the differences which occur between his results and mine. Thus, in the table of the heights of the boys of eleven years of age already given (page 18), we find a variation of fourteen groups of an inch each; that is to say, some of the boys are 13 inches taller than others of the same age; and a much wider difference occurs as the age advances; so that it could be by accident only that ten of the groups could determine the true mean boy: there would not be one boy to represent

each group which might occur in 430 individuals, and the three omitted groups would influence the position of the mean, especially if the mean was a mere mathematical one. The accuracy, therefore, of M. Quetelet's figures depends on his ability to select 'properly constituted' models for measurement; and as such selections might vary with the experience of observers, the results of two or more observers could hardly be identical, or their typical forms or standards the same. If the typical form could be eliminated in this manner, the limits and the grouping of the variations could not, which are matters of equal practical importance. The method which I have adopted of accumulating a large number of measurements for each age is the only one which will meet all these requirements.

The human type is not so fixed and persistent in its characteristics as not to be liable to yield to the influence of constant causes. The Chinese will always be shorter in stature than Europeans, but still the law of accidental causes will not be less constant in its operation; the only result will be that in each race the variations in height, &c., will take place round a greater or smaller mean, and they will in both instances be influenced by climate, food, occupation, &c. These regular oscillations around the same type are observable from the very instant of

birth, only their limits become more confined the nearer we approach that period; as if Nature, acting alone, was loth to deviate from its type, and as if the greatness of the variations arose chiefly from the modifying influences of man.\* It is for the purpose of fixing the typical forms of man for each age as they exist *at present* in this country, and for ascertaining the range and nature of the deviations from those forms, and the modifying influences producing the deviations, that my present work is written. It is necessary to bear in mind that the typical form eliminated by this method is not necessarily the most perfect form of man, but represents the equilibrium, as it were, of many contending forces which may be disturbed by the future predominance of any one of them; hence the typical form is not the same for the working and the non-working man, for the man living in towns and the man living in the rural districts. In this way each group above and below the mean is typical of the predominance of some modifying cause,

\* It is in man and domesticated animals that the widest range of variation is observed. In wild animals, and especially in birds and insects, the individuals of the same species appear to differ little in size; and the naturalist often relies on the measurements of the length of the body, the wings, &c., as important distinctive marks of varieties. Naturalists are, however, accustomed to speak of 'fine specimens,' which shows that there are important accidental variations from the mean.

if we could but distinguish it. Many of the modifying causes are obscure and very complicated; but others are easily recognised, and are deserving of careful investigation.

While adopting M. Quetelet's method of studying the proportions of the body as the best, and indeed the only, scientific method yet propounded, we cannot accept his tables with equal confidence, as they are based on too small a number of observations; and, moreover, the measurements, being those of Belgians, are not applicable to the inhabitants of this country. In my tables in the eighth volume of the *St. George's Hospital Reports*, I have shown the differences which exist between M. Quetelet's averages and my own, which are based on a much larger number of observations. Professor Bowditch, in his work *On the Growth of Children in America*, has also called attention to an interesting and important fact, which appears to have escaped M. Quetelet's notice, that at the age of 13 and 14 years girls are taller and heavier than boys of the corresponding ages; and I have shown that the same is true in this country. M. Quetelet represents the two sexes as of the same size at 12, but at all other ages the female is smaller than the male. M. Quetelet found the inhabitants of towns taller than those living in rural districts, the reverse of the order which exists in this country.



## CHAPTER II.

### APPARATUS.

THE apparatus for making measurements of the body and for ascertaining the weight and strength are few in number, and, with the exception of the callipers and the dynamometer, are easily accessible in some of the common forms in use for commercial purposes. But as I am anxious to encourage anthropometry as a systematic study, not only among scientific men and the members of the medical profession, but among the managers of gymnasiums and schools, I shall describe the instruments which I have found most convenient for the purpose, many of which have been constructed according to my instructions.

*The unit of measurement is the English inch divided into tenths and not eighths, as in the ordinary rules in use in this country. The decimal system of notation occupies less space in the chart, is most easily written down, and is best adapted for subsequent arithmetical and statistical manipulation.*

On the hand-rule, the measuring-rod, and the measuring-tape which I am accustomed to use, the

metrical and English system of measures are placed in juxtaposition, and both can be read off at the same time with equal facility. This arrangement saves a great deal of troublesome calculation, and helps to familiarise the mind and the eye with the relative values of the two systems, and enables the observer to register his measurements in metrical and English values in parallel columns. Where calculations have to be made to reduce the one system to the other, the following table will be useful:

Measures of length.	In English inches.	In English feet (=12 inches).	In English yards (=3 feet).
Millimètre . . .	0·04	—	—
Centimètre . . .	0·39	0·03	0·01
Décimètre . . .	3·93	0·32	0·10
Mètre . . . . .	39·37	3·28	1·09
1 inch equals 2·54 centimètres. 1 foot „ 3·04 décimètres. 1 yard „ 0·91 mètres.			

N.B. If the measurements be taken in inches and *eighths*, the fractions must be reduced to the decimal form, thus:

Eighths of an inch reduced to tenths.

1	inch equals	1·00
$\frac{7}{8}$	„ „	0·87
$\frac{3}{4} = \frac{6}{8}$	„ „	0·75
$\frac{5}{8}$	„ „	0·62
$\frac{1}{2} = \frac{4}{8}$	„ „	0·50
$\frac{3}{8}$	„ „	0·37
$\frac{1}{4} = \frac{2}{8}$	„ „	0·25
$\frac{1}{8}$	„ „	0·12

*The unit of weight and strength is the English pound avoirdupois (16 ounces). I am not aware of the existence of any English weighing-machines adapted for the use of metrical weights of which the kilogramme=2·2 pounds English would form the unit, as there is at present no demand for them; but if they were required the mechanical difficulty could be easily overcome, and the employment of two sets of weights of different shapes would prevent confusion or mistake. The following table will be useful for reducing the English to the metrical weights:*

Measures of weight.	In English ounces, avoird.	In English pounds (=16 ounces), avoird.	In English stones (=14 lbs.), avoird.
Gramme . . . .	0·03	—	—
Décagramme . .	0·35	0·02	0·001
Hectogramme . .	3·53	0·22	0·015
Kilogramme . .	35·27	2·20	0·157
1 lb. avoird. equals 0·45 kilogrammes.			
1 stone „ „ 6·35 „			

We are indebted to Mr. John Bellows of Gloucester for an admirable contrivance for reducing the English to the metrical, and the metrical to the English system of weights and measures. It consists of a series of circular cards placed one above the other, and diminishing in size from the bottom to the top, the whole of them working on a common centre. Round the margins of the cards are printed the weights or measures of the two systems, and by turn-

ing the cards so as to bring the required values opposite to each other the result can be read off by a simple process of addition. For my own method, see the remarks on the *measuring-staff*.

The *hand-rule* is an ordinary boxwood measure one mètre in length (39·37 inches English), with four joints to allow of its being folded up in one direction. It is graduated into English inches and tenth of inches, and the mètre is subdivided into décimètres, centimètres, and millimètres. The two scales are placed in direct apposition to show their relative values, but are easily distinguished by the difference in the size of the figures employed for marking the two systems. The hand-rule, besides being the standard of reference, has other uses which will be mentioned below.

The *measuring-staff*, for taking the various heights of the body, is a light square rod of lance-wood two mètres in length, with a joint in the middle for convenience of carriage or storing away when not required for use. Its total length is 78·74 inches English, or rather more than 6 feet 6 inches. It is graduated on the face into feet, inches, and tenths of inches; and, to save the trouble of calculation and to prevent mistakes being made, the whole of the inches are marked in succession, from 1 at the bottom to 78 at the top. On one side of the staff the





Diagram of a portion of the scales on the front, side, and back of the measuring-rod, showing the relative value of English inches and centimetres, and of English pounds avoirdupois and kilogrammes. The two divisions—*inches* and *centimetres*—represent a portion of the *hand-rule* actual size.

metrical system is given, the smaller divisions of which meet at the same angle as the smaller divisions of the English measures, to show their relative values. On the back of the rod a scale is given, which, in conjunction with the centimètre scale, is intended to facilitate the conversion of English pounds weight into kilogrammes, or the reverse. For this purpose the 200 centimètres of the length scale are made arbitrarily to represent 200 pounds weight, and the divisions on the back of the rod 91 kilogrammes, the slight fractional difference being disregarded as of no practical importance. A brass loop is made to slide stiffly up and down the rod by means of a spring to facilitate reference to the scales on different surfaces of the rod, and into the loop the hand-rule can be fitted to form the cross-bar when heights are being taken.

The *measuring-tape* should be two mètres in length, and graduated in the same manner as the hand-rule. As the graduations on measuring-tapes are frequently incorrect, and as the marks influence the mind and hand in determining measurements, especially in taking circumferences of soft parts, it is better to dispense with them and use a perfectly plain tape, and read off the distance on the hand-rule. This is absolutely necessary when great accuracy is desired, as the most practised observer cannot

resist the habit of fixing on the larger and the *even* divisions, rather than on the smaller and *odd* divisions of a scale.

The *callipers* which I use are made of boxwood, and are of a pattern sometimes used by artists. They are 15 inches in length, and are available for measuring from a fifth of an inch to 20 or more inches. They are furnished with a light brass arc, carrying a scale of English inches on one side and of centimètres on the other.



Boxwood Callipers, with brass scale of inches (Eng.) and centimètres.

The *weighing-machine* which appears to me the most convenient, accurate, and suitable alike for public institutions and private consulting-rooms, is made by Messrs. Young & Son of Cranbourne-street, and is designed specially for weighing human beings.\*

\* This is a very old-fashioned machine. In the Conservatoire des Arts et Métiers in Paris there are five worn-out instruments of this kind, all of which are evidently of English manufacture, as the standards are graduated in English feet and inches, and one of them bears the name and address of the maker, Henry Kettle, St. Paul's Churchyard—a name which has disappeared from the *London Directory* of the present day. In the museum catalogue they are entered thus: *R. f. 46 Romaine, dite de Sanctorius*. As far as I can judge, the alterations made in the machine by Messrs.

A very good machine, of the ordinary commercial shape, is sold by Mr. Hawksley of Oxford-street,\* which has the advantage of being much less expensive than Messrs. Young's instrument. The numerous weighing-machines to be found in railway stations and other public places are sufficiently accurate when the weight is taken in the clothes, but they are too cumbersome and expensive for private use. In Messrs. Young's and Mr. Hawksley's machines the weight is taken in the upright position, and measuring-rods are attached, so that the height also can be taken while the person under examination

---

Young add greatly to the expense without increasing its usefulness. The objections to the machine are the use of the old-fashioned scale-pan for the weights instead of the simple beam and suspender adopted in modern weighing-machines, and the demand for additional space made by the pan projecting behind, instead of at the side of, the instrument.

\* This machine is approved of by Dr. Fergus of Marlborough College, who used it for taking the weight of the boys; and it was used in the various schools in Boston, U.S.A., and neighbourhood, for ascertaining the weights of about 24,500 children, referred to by Dr. H. P. Bowditch in his work on the growth of children in the United States of America.






is still on the scales, an arrangement which saves much time and trouble when a large number of individuals are being examined.

The dynamometer, to be immediately described, can be used as a weighing-machine, if it is suspended to a beam or try-pode by one handle, and a stirrup or bar for the person who is to be weighed be attached to the other. The makers of the dynamometer have fitted it with the necessary appliances for use as a weighing-machine at my suggestion.

A very convenient *dynamometer*, for testing the strength of the hands (grasping), the arms (pulling),




THE DYNAMOMETER.

and the back (lifting), is made by Mr. Coxeter of Grafton-street East, W.C. It consists of a bar of steel coiled on itself in the centre, the ends being left free

and placed parallel to each other, to act as levers. To these levers are attached handles, which can be adjusted according to the test it is proposed to apply. In front of the coil is placed a graduated scale of pounds and kilogrammes, with an index and registering-pointer. The action of the instrument is very simple, and it is available for testing the strength of children as well as adults.\*

The callipers, hand-rule, and measuring-tape are important aids to the physician and surgeon for the diagnosis of diseases, accidents, and deformities of the body, and should be found in the wards of every hospital and private consulting-room. I have had my instruments fitted into a light case for the sake of portability, and for preserving them from injury and preventing misplacement or loss. This case I call my 'physical examination case.' The dynamometer, fitted also as a weighing-machine, is packed into a small convenient case; so that the two cases, with a few loose sheets of the anthropometrical chart, form a complete 'outfit' for the anthropometrical student, a substitute for the measuring-rod being easily extemporised.

\* This instrument is not adapted, however, for testing the *lifting power with both hands of very strong persons*, as the delicacy necessary for testing the grasping power would be lost in an instrument suitable for that purpose.



*The Dynamometer arranged as a Weighing-machine.*—Men and boys may be weighed by simply holding the cross-bar with the hands'; children and girls by putting the arms through the padded straps, or by reversing the bar and straps, and sitting on the bar as in a swing. The weight of the bar and the straps must be deducted from the total weight, as it cannot be allowed for in the machine without interfering with its use in determining the strength.

## CHAPTER III.

### THE ANTHROPOMETRICAL CHART.

THIS chart consists of three portions. The first, on the left, contains an outline *diagram* of the human form; the second or central portion, consisting of a series of vertical columns, is the *register*; and the third portion, on the right of the sheet, is the *chart* properly so called. The diagram of the body and the columns of the register are crossed by numerous horizontal lines, which indicate on the diagram the parts of the body to be measured, and on the register the exact place where the measurements are to be written down. The first and second columns of the register contain the names of the parts to be measured, so that the diagram may be dispensed with by persons who are familiar with the anatomy of the external form of the body. To economise space, the heights of various parts of the body from the sole of the foot, and the distances downwards from the vertex, and from the trochanter, are written in ~~the~~ the same column of the register, the heights *above* and the distances *below* the horizontal lines corresponding to the parts measured. The transverse and the

antero-posterior diameters of the body are also written in the same column, the former *above* and the latter *below* the lines. The fine hair-line numbers in the columns of the register refer to the *Systematic Table of Measurements* (p. 47), where a description of the proper method of making each measurement is given. These reference numbers are engraved in hair-lines, in order that they may be written upon and obliterated by the actual measurements, as they also mark the exact spot where the entries are to be made. The chart proper is ruled into divisions of inches and tenths of inches, for tracing, by the graphic method, the variations which take place in the size, weight, and strength of the person examined. The scale is indicated by the three columns on the left, one division of the tenth of an inch being equal to one inch in measurements of size, to two pounds in weight, and of five pounds in measurement of strength. The perpendicular lines of the chart represent the dates at which the observations are made, which should be at regular intervals of time: thus, one division of a tenth of an inch should correspond to one day, one week, one month, or one year. Another use of the chart is to lay down a plan or outline diagram of the body, either in front or in profile, from the measurements given in the register, and to make this the starting-point

for marking subsequent variations which may result from growth, training, or disease. (See the Chart-tracings, Nos. 1, 2, 3). Observers who are familiar with the graphic method of treating statistics may dispense altogether with the diagram of the body and the register, and mark on the chart itself the measurements at the time they are made on the body. This remark applies more especially to the recording of clinical observations, and the sheet may be divided in the middle, leaving the column containing the 'General Remarks' attached to the chart.\*

It will be understood that, although the diagram represents the figure of a young man, the register and chart are intended for recording the physical proportions of both sexes at all ages, as the parts of the body to be measured are the same in all.† Some of the measurements are of more importance than others; and these are indicated by the horizontal black lines on the diagram, while those of secondary importance are indicated by dotted lines. When the chart is used for special objects of study, the student

\* 'Statistical results are now frequently expressed by graphic representations, a certain space drawn to a scale representing a number. The most simple plan is that of intersecting horizontal and vertical lines. . . . Such graphic representations are most useful, and allow the mind to seize more easily than by rows of figures the connection between two conditions and events.' *A Manual of Practical Hygiene*, by E. A. Parkes, M.D., p. 514.

† When used for women or children the figure can be torn off.

may make use of those measurements only which refer to his subject, or substitute others which he may deem more suitable; but it is desirable in all cases where it is possible, to make a thorough examination of each person, and afterwards to record on the chart the variations which are more especially worthy of notice.

THE GENERAL REMARKS refer to the personal identity of the individual examined (name, age, sex, complexion), his nationality or racial origin (Celtic, Teutonic, Scandinavian), the circumstances which surround him and modify his bodily conformation (occupation, nurture, sanitary surroundings), and the results of those modifying influences (imperfect development, deformity, disease).

The *date*, *name*, *sex*, and *age* need no explanations. The complexion or temperament is determined by the *colour of the hair*, which is distinguished as *fair*, *red*, *brown*, *dark brown*, *black*; and by the *colour of the eyes*, which may be described as *light blue*, *dark blue*, *gray*, *brown* (or *hazel*).

*Racial Origin.* The terms English, Scotch, Irish, and Welsh very imperfectly express the racial origin and national peculiarities of different portions of inhabitants of the British Isles and their descendants in America and Australia, and should be avoided when the physical conformation and development of

the body are under consideration. The original elements of our British race are becoming so intimately mixed by intermarriage, by easy communication between one part of the country to another, by railways, and by the fluctuations of commercial enterprise, that it becomes every day more and more difficult to distinguish them. But the following table, which indicates the parts of the British Isles which were originally occupied by the Celtic, Teutonic, and Scandinavian races, will in a measure assist the student to distinguish their descendants:

#### CELTIC RACES.

Highlands of Scotland.	West and southern por-
North and South Wales.	tions of Ireland.
Devon and Cornwall.	

#### SCANDINAVIAN RACES.

Shetland.	Northumberland.
Eastern Lowlands of Scot-	Durham.
land.	Yorkshire.
Dumfriesshire.	Lincolnshire.
Cumberland.	Nottinghamshire.
Westmoreland.	Leicestershire.

#### TEUTONIC RACES.

The English counties not enumerated in the above lists. The inhabitants of the English counties border-



ing on Wales, and on the north and east coast of Ireland, are a mixture of these races.\*

*The occupation* of an individual, or, if a child, of its parents, explains not only the direct effect of physical or mental work on the constitution of the body, but the kind of nurture and sanitary surroundings to which he may have been subjected. The following classification has been arranged to show all these influences, beginning with the most favoured and ending with the least favoured class of the population of this country. The statistics are taken from the census returns of 1871, and the percentages include the children distributed over all the classes.

*Classification of Occupations in England, according to their Social Position and Sanitary Influences.*

Upper and Professional Classes.	CLASS I.		328,270	4.46
	(a) <i>Outdoor Occupations.</i>			
	Persons of Rank.			
	Officers of Army and Navy.			
	Clergymen, Doctors.			
	(b) <i>Indoor Occupations.</i>			
Commercial Classes.	Lawyers.		762,014	10.36
	Civil Servants.			
	Bankers, Merchants.			
	CLASS II.			
	<i>Indoor Occupations.</i>			
	Clerks.			
	Shopkeepers.			

\* See Dr. Beddoe, 'On the Stature and Bulk of Man in the

Labouring Classes.	CLASS III. <i>Selected Occupations.</i>		
	Soldiers.	439,377	5.97
	Policemen.		
	Domestic Servants, &c.		
	CLASS IV. <i>Outdoor Labourers.</i>		
	Agriculture, Roads, Quarries, Railways, &c.	2,383,799	32.41
Industrial Classes.	CLASS V. <i>Seafaring Occupations.</i>		
	Sailors.	232,784	3.16
	Fishermen, Bargemen.		
	CLASS VI. <i>Underground Occupations.</i>		
	Miners, Coal and Mineral.	435,558	5.92
	CLASS VII. <i>Indoor Occupations.</i>		
	Artisans.	1,971,295	26.82
	CLASS VIII. <i>Indoor Occupations.</i>		
	Factory Operatives.	801,536	10.90
	Sedentary Trades, Tailors, &c.		
	Children of no Occupation. . .	3,704,301	
Total Male Population. . . . .		11,058,954	100.00

The *hereditary diseases* which affect the conformation of the body are scrofula, syphilis, rickets, idiotcy, &c. The *congenital deformities* are the absence or imperfect development of organs, club-foot, supernumerary members, &c.; and the *acquired de-*

British Isles,' *Memoirs of the Anthropological Society of London*, vol. iii. p. 166, 1870.

*formities* are flat-foot, knock-knee, weak ankles, curvature of the spine, &c. Other circumstances which may influence the development of the body should be recorded, such as exposure to different climates, peculiar modes of living, unusual physical proportions of parents, left-handedness, &c.

NOTE.—Naturalists, travellers, missionaries, and others engaged in anthropometrical inquiries outside the British race and its offshoots will find in the following classification of the races of mankind the principal characteristics deserving of record :

‘The different races of mankind are divisible into two primary divisions ; the *Ulotrichi*, with crisp or woolly hair, and the *Leiotrichi*, with smooth hair.

‘*a.* The colour of the *Ulotrichi* varies from yellow-brown to the darkest hue known among men. The hair and eyes are normally dark, and, with only a few exceptions (among the Andaman Islanders), they are *dolichocephali*. The negroes and bushmen of ultra-Sahara Africa, and the Negritos of the Malay Peninsula and Archipelago, and of the Papuan Islands, are the members of this *Negroid* stock.

‘*b.* The *Leiotrichi* are divisible into :

‘1. The *Australoid* group, with dark skins, hair, and eyes, wavy black hair, and eminently long prognathous skulls, with well-developed brow ridges, who are found in Australia and in the Dekhan.

‘2. The *Mongoloid* group, with, for the most part, yellowish-brown or reddish-brown skins, and dark eyes ; the hair being long, black, and straight. Their skulls range between the extremes of dolichocephaly and those of brachycephaly. These are the Mongol, Tibetan, Chinese, Polynesian, Esquimaux, and American races.

‘3. The *Xanthochroic* group, with pale skins, blue eyes, and abundant fair hair. Their skulls, like those of the Mongoloid group, range between the extremes of dolichocephaly and brachycephaly. The Slavonians, Teutons, Scandinavians, and the fair Celtic-speaking people are the chief representatives of this division ; but they extend into North Africa and Western Asia.

‘4. The dark whites, or *Melanochroi* ; pale-complexioned people, with dark hair and eyes, and generally long, but sometimes broad, skulls. These are the Iberians and “black Celts” of Western Europe, and the dark-complexioned white people of the shores of the Mediterranean, Western Asia, and Persia.’ *The Anatomy of Vertebrated Animals*, by T. H. Huxley, F.R.S., 1874, p. 496.

## CHAPTER IV.

### A SYSTEMATIC TABLE OF MEASUREMENTS OF THE HUMAN BODY.

#### *Preliminary Remarks.*

WHEN we endeavour to describe with accuracy the proportions of the external form of the body, we meet with difficulties in defining the various points from which measurements can be taken. We depend chiefly on the prominent points of the bony framework, and rely as little as possible on the soft parts; a few points of which, however, are sufficiently definite and circumscribed to be available for our purpose, such as the nipples and the navel. Whether we make measurements of the human body or of inanimate things, errors of observation are liable to occur; but it is well known that all such errors are subject to a definite law—the law of the frequency of error—and they can be eliminated by making several measurements, and deducing from them an average which will be the nearest approach to the actual measurement required.

It must be understood that the following measurements refer to the naked body. The heights must be

taken without the shoes, and the diameters and circumferences must be made on the skin itself. Too much importance cannot be attached to this rule, as its non-observance will render the labour worthless and the results misleading, as has been the case with a large amount of the statistics of measurements of the body which have been already published. If, however, the rule cannot be strictly observed, the height of the shoes and the thickness of the dress should be ascertained and deducted from the measurement before it is entered in the register, in order to avoid future error or confusion.\*

In recording measurements on the anthropometrical chart, the following instructions must be strictly carried out:

The reference numbers of the Systematic Table of Measurements (to be given immediately) are printed in fine hair-lines in columns 3, 4, 5, 8, and 9 of the register, and indicate the places where the actual measurements are to be written down (*i.e.* the mea-

\* The absurdity and false modesty which requires that the whole body, with the exception of the hands and face, should be mysteriously hidden by a mass of tight-fitting clothes, is much to be regretted. Children especially ought to have more freedom in this respect, and should be systematically examined without their clothes by competent persons, who can judge of their physical development, and detect at the earliest moment any departure from the normal state.

surements are to be written *upon* the reference numbers).

Measurements of *height*, from the sole of the foot, are to be written *above* the lines corresponding to the parts measured; and *distances* from the vertex, from the acromion and the trochanter, *below* the lines.

*Transverse diameters* are written *above*, and *antero-posterior diameters* are written *below*, the lines corresponding with the parts measured.

*Circumferences* are written *above* the lines, and the *breadths* of the hand and foot *below* the lines, corresponding with the parts measured.

All the measurements must be given in English inches and tenths of inches (*i. e.* in decimals), and the weights and strengths in English pounds avoirdupois (16 ounces), unless the metrical system be adopted.

#### I. HEIGHT FROM THE SOLE OF THE FOOT (*i. e.* from the Ground).

For taking the heights of various parts of the body from the sole of the foot the person under examination must stand in the upright position without shoes, and the measurements be made with the measuring-staff. It is necessary to observe this rule, as the total height of the body is greater in the recumbent than the upright position. In the case of infants the recumbent position must necessarily be assumed, and

the measurements may then be more easily made with the callipers.

1. Height from the sole of the foot to the perineum or fork.

For the measurements of the lower limb see Nos. 41, 42, and 43.

2. Height from the sole of the foot to the pubes.
3. Height from the sole of the foot to the abdomen, on a line with the anterior superior spines of the iliac or haunch bones.
4. Height from the sole of the foot to the navel.
5. Height from the sole of the foot to the lower end of the sternum or breast-bone (enciform cartilage).
6. Height from the sole of the foot to the top of the sternum or breast-bone (sterno-clavicular articulation).
7. Height from the sole of the foot to the chin, the head being held in a perfectly horizontal position.
8. Height from the sole of the foot to the vertex (total height).

## II. DISTANCES FROM THE VERTEX.

The measurements of the face are made by folding the hand-rule to a right angle, and resting one limb on the vertex and the other on the tip of the nose.

The perpendicular distances are read off by noting on the rule where the horizontal lines corresponding to Nos. 9, 10, 11, 12, and 13 intersect it.




Fig. I.

9. Perpendicular distance from the vertex to the beginning of the growth of the hair on the forehead.
10. Perpendicular distance from the vertex to the frontal sinuses (*i.e.* the eyebrows and the root of the nose).
11. Perpendicular distance from the vertex to the basis of the nose.
12. Perpendicular distance from the vertex to the mouth.
13. Perpendicular distance from the vertex to the point of the chin.

E

LANE LIBRARY



The distances from the vertex to various parts of the trunk need not be enumerated here, as they are between the same parts as the heights already given, but measured in the reverse direction. If they are made, they must be written *below* the lines corresponding to the respective heights and parts measured.

### III. DIAMETERS AND CIRCUMFERENCES OF THE HEAD AND TRUNK.

All the diameters of the body are measured with callipers, and the circumferences with the measuring or the plain tape; the pressure made when using the instruments should not be so great as to produce indentation of the skin. As the diameters and circumferences of the trunk vary with the state of the respiration, the measurements should be taken when the chest is empty and at rest. This position is most easily secured by making the person under examination count in a loud voice from one to ten, immediately before determining the measurements, the arms being allowed to hang loosely by the sides of the body.

The accompanying diagrams from Kirke's *Handbook of Physiology*, page 208 (after Hutchinson), will convey an idea of the changes which take place in the shape and size of the trunk under the influence of natural and forced respiration. The former is repre-

YMA 3341 1754

sented by the thick black lines and the latter by the dotted lines.




Fig. II.

14. Transverse diameter of the head above the ears on a line with the frontal sinuses. (Fig. I. No. 10.)
15. Antero-posterior diameter of the head from the frontal sinuses to the occipital protuberance. (Fig. I. No. 10.)
16. Circumference of the head on a line with the frontal sinuses (eyebrows). (Fig. I. No. 10.)
17. Transverse diameter of the face below the ears at the articulations of the lower jaw. (Fig. I. No. 11.)
- 17A. Antero-posterior diameter from the base of the

nose to the back of the neck (spine of first vertebra). (Fig. I. No. 11.)

18. Circumference of face and head on a line with the base of the nose. (Fig. I. No. 11.)
19. Transverse and antero-posterior diameters, and the circumference of the neck on a line with the chin. (Fig. I. No. 13.)
20. The same measurements of the neck at the root of the neck where the clavicles meet.
21. Transverse diameter of the shoulders between the acromion processes.
22. Circumference of the shoulders at the same place.
23. Transverse diameter of the chest at the axillæ.  
Circumference at the same part (the chest empty).
24. Transverse diameter of the empty chest on a line with the nipples.
25. Antero-posterior diameter of the empty chest on a line with the nipples.  
N.B. The distance between the nipples may be written on the *diagram* between the points measured.
26. Circumference of the empty chest on a line with the nipples.

NOTE. As the principal object of making anthropometrical observations is to ascertain the size of the body and not its fitness for any special functions, the

above measurement is devised for showing the size, and not the breathing capacity, of the chest. The measurement is the one in use in our army and navy for the examination of recruits, and is thus effected: A measuring-tape is placed quite horizontally round the chest, the lower edge of the tape touching the upper part of the nipple and including the lower portions of the scapula behind, the arms hanging loosely by the sides. The recruit is made to count from one to ten to prevent undue inflation of the chest, and the length shown on the tape in inches is the correct chest-girth.

- 26 +. The difference between the empty and the full chest-girth; to be written opposite the plus (+) sign above No. 26 (expansibility of chest).
- 27. Transverse antero-posterior diameters and circumference of the empty chest at the lower end of the sternum (breast-bone).
- 28. Transverse antero-posterior diameters and circumference at the waist.
- 29. Transverse and antero-posterior diameters at the crest of the pelvis (haunches).
- 30. Circumference at the same part.
- 31. Transverse diameter at the trochanters (hip-bones).
- 31A. Antero-posterior diameter at the pubes.
- 32. Circumference at the trochanters (the hips).

## IV. THE UPPER AND LOWER LIMBS.

The measurements of the upper limb are made when it is extended horizontally on a line with the shoulder,\* and the lower limb either in the upright or recumbent position. The points of departure are, for the upper limb, the acromion prominence of the shoulder, and, for the lower limb, the trochanter. The dotted lines on the right arm and leg of the diagram indicate the direction and the points between which measurements are to be made. Circumferences of muscles, such as the biceps, must not be made immediately after they have been exercised, as they are then unduly distended with blood.

a. *Upper Limb.*

33. Length of the two arms extended horizontally without forced extension, measured from the tip of one middle finger to that of the other.
34. Length of the arm from the acromion (tip of the shoulder) to the external condyle of the humerus at the elbow-joint.
35. Length of the fore-arm from the external condyle of the humerus to the styloid process of the radius at the root of the thumb.

\* This position of the arm has not been represented in the anthropometrical chart, on account of the large addition to the size of the chart it would render necessary.

36. Length of the hand from the styloid process of the radius, across the back of the hand, to the metacarpal joint (knuckle) of the middle finger.
37. Length of the middle finger from the knuckle (metacarpal joint) to the tip of the finger.
38. Circumference of the arm at the middle of the biceps muscle.
39. Circumference of the fore-arm at the thickest part.
40. Circumference of the hand at the metacarpal joints (knuckles).
- 40A. Breadth of the hand at the same part (to be written below the line).

*b. Lower Limb.*

41. Length of the thigh from the trochanter to the lower edge of the patella (knee-cap).
42. Length of the leg from the lower edge of the patella to the internal malleolus (the inner ankle-bone).
43. Length (*height*) of the foot from the internal malleolus (ankle-bone) to the sole of the foot (*i.e.* to the ground).
44. Length of the foot from the heel along the inside of the foot to the top of the metatarsal joint (the ball) of the great toe.

45. Length of the great toe from the top of the metatarsal joint to the tip of the toe.
46. Circumference of the foot at the metatarsal joints.
- 46A. Breadth of the foot at the same part (to be written below the line).
47. Circumference of the leg above the ankle-joint.
48. Circumference of the leg at the calf of the leg.
49. Circumference of the knee.
50. Circumference of the thigh at the middle.
51. Circumference of the thigh at the fork.

#### V. WEIGHT OF THE BODY.

The weight of the body should be taken without the clothes. When this is not possible, the *whole of the indoor dress with the shoes should be included*, and the weight of the dress should be recorded separately. Food and exercise will make the weight of the body vary to the extent of two or three pounds, and the weight, therefore, should be taken at the same time of the day. The best time is in the morning before breakfast.

52. Weight of the body in pounds avoirdupois taken without the clothes.
53. Weight of the body in pounds avd. taken in the ordinary indoor dress, including the shoes.

NOTE. The weight of the clothes may be

written in either of the numbers 52 or 53 left vacant.

## VI. STRENGTH OF THE BODY.

In determining the strength by the dynamometer it is advisable to make three trials at intervals of a few minutes and record the average. It is necessary that the instrument should be held or placed in such positions as to prevent the weight of the body interfering with the trials of strength.

### a. *Both Arms.*

- 54. Strength of the arms in antagonism—'pulling'—the instrument being placed in front of the body.
- 55. Strength of the back—'lifting' with both hands—the instrument being fastened to the ground or by a strap to the feet.

### b. *Right Hand.*

- 56. Strength of the right hand shown by 'grasping' the handles of the instrument together.
- 57. Strength of the right arm shown by 'pulling,' the instrument being held in front of the body to prevent the weight being thrown in.
- 58. Strength of the right arm and back shown by 'lifting,' the instrument being in the position of No. 55.



*c. Left Hand.*

59. Strength of the left hand shown by 'grasping:'  
see No. 56.
60. Strength of the left arm shown by 'pulling:' see  
No. 57.
61. Strength of the left arm and back shown by 'lift-  
ing:' see No. 58.

## CHAPTER V.

### RELATIVE PROPORTIONS OF THE BODY.

THE measurements described in the foregoing table are to be made *directly*, and show the *absolute* values for each part of the body. But it is also important to ascertain the relative proportions, and the relation which each part bears to the whole. This is done by considering the total height as constant, and equal to 1, or to 100, and calculating what portion of the total height each part of the body forms; and the upper and lower limbs are to be treated in a similar manner. The first, second, sixth, and seventh columns of the Anthropometrical Chart are available for recording these relative proportions of the body, which can be calculated from the *direct* measurements given in the other columns of the register. In representing relative proportions by tracings on the chart the curves will be the reverse of those of the *direct* measurements, and will bend downwards instead of upwards, as shown in the Diagrams II. and III.

## RELATION OF WEIGHT TO HEIGHT.

In order to ascertain with accuracy the relative weight and height of the human body, it is necessary to determine for each age and in each group of measurements the average weight corresponding to each height. The following table, published by Mr. G. C. Steet,\* showing the relation of the weight to the height of a large number of boys at the Telegraph Department of the General Post Office, London, will illustrate this mode of procedure much better than a verbal description. I have added to the table a column of averages of the weights at each height in inches, to show how the weight varies with the age in boys who possess the same stature. I have also altered the arrangement of the table to make it conform with my other tables, but the results have not been interfered with.

\* 'The Development and Growth of Boys between Thirteen and Twenty Years,' by G. Carrick Steet, F.R.C.S.,—*St. George's Hospital Reports*, 1874-6, p. 52.

*Table showing the Relation of Weight to Height of Boys between the Ages of 13 and 20 Years.*

[illegible]

This method of ascertaining the relation of the weight to the height of the body involves a large amount of labour, and has the further disadvantage which results from the employment of *averages* instead of *means* (see post, page 69), inasmuch as exceptional cases influence averages and produce irregularities in the series of results, while means, by disregarding exceptional cases, produce regularity and uniformity in the series.

Another method of finding the relation of the weight to the height is to divide the weight in pounds by the height in inches, as I have done in Table IV., in the column headed 'Ratio of height to weight and chest-girth.' This method shows the weight for each inch of stature for each age from 10 to 30 years. A typical boy of 10 years of age weighs 1·252 lbs. for each inch of his height, and one of 23 or 24 years 2·203 lbs. By multiplying these figures by the total height, we obtain the typical weight for their respective ages.

A third method of arriving at the same end is to arrange the heights and weights for each age opposite each other in groups which will bring them within the same limits, as I have done in the following table of the heights and weights of 430 town and country schoolboys, of the age from 11 to 12 years. The statistics were derived from various sources, and

the measurements and weights were taken by several independent observers. The table shows that for each inch in height there is an increase of weight of 3·5 lbs.\*

*Table showing the Relation of the Heights and the Weights of 430 Schoolboys between the Age of 11 and 12 Years.*

	Heights in groups of 2 inches (without shoes).	No. of boys.	Weight in groups of 7 lbs. (including clothes).	No. of boys.	The diagrams are drawn to a scale. The black lines represent the weights, and the dotted lines the heights.
	60 to 62	3	98 to 105	4	
	58 " 60	16	91 " 98	15	
	56 " 58	58	84 " 91	62	
Mean	54 " 56	133	77 " 84	133	
	52 " 54	137	70 " 77	129	
	50 " 52	64	63 " 70	77	
	48 " 50	16	56 " 63	10	
	46 " 48	3	49 " 56	?	
		430		430	

This method can also be applied to the average and mean heights and weights for each age, for the purpose of ascertaining their relation to each other; and although the result is not strictly accurate, it is an approximation sufficiently near the truth for all practical purposes. The mean heights

\* See the *Physical Requirements of Factory Children* (London, 1876). 'The Growth of Children,' by H. P. Bowditch, M.D., *Eighth Annual Report State Board of Health, Massachusetts* (Boston, U.S., 1877).

and weights corresponding to each age should be arranged in parallel columns, and the weights corresponding to each even inch determined by interpolation. The tracing on Diagram III., showing the relation of the weights to the heights given in Tables I. and II., has been constructed on this plan, and the following figures will show how it was effected:

Age.	Observation.		Interpolation. Weight.	Rate of increase for each even inch of height.	Age.	Observation.		Interpolation. Weight.	Rate of increase for each even inch of height.
	Mean Height.	Mean Weight.				Mean Height.	Mean Weight.		
	inches.	lbs.	lbs.	lbs.		inches.	lbs.	lbs.	lbs.
10	53	67	67			62	110	108	4
	53·5		69			62·5		105	
	54		71	4		63		107	4
11	54·5	73	73		15	63·5		110	
	55		75	4		64		113	6
	55·5		77			64·5		116	
12	56	80	79	4		65	126	119	6
	56·5		81			65·5		122	
	57		83	4		66		125	6
13	57·5	88	85		16	66·5		129	
	58		87	4		67		132	7
	58·5		89			67·5		135	
14	59	98	91	4	17	68	140	140	8
	59·5		93		18	68·5	146	146	
	60		95	4	19	69	148	148	8
14	60·5	98	97		20	69	150	150	
	61		99	4	21	69	152	152	
	61·5		101		25 to 30	69	154	154	6

Thus, for each inch in height the weight is increased 4 lbs. from 10 to 15 years; 6 lbs. from 15 to 16; 8 lbs. from 16 to 19; and from 19 to 30 years of age there is 6 lbs. added to the weight, while the height remains stationary.

## CHAPTER VI.

### STATISTICAL TABLES.

IN combining into statistical tables the measurements and observations obtained in the manner I have indicated in the foregoing pages, it is of the utmost importance that a uniform plan should be adopted, to admit of ready comparison of the results of different observers. Statistical tables, moreover, should be constructed with a completeness of detail which will preclude the possibility of doubt, or of mistakes being made even by persons imperfectly acquainted with the subjects to which they refer. They should also be diagrammatic, in order that the eye may take in at a glance the relation of the facts which they are intended to portray.

I submit the following original tables of the height, weight, and chest-girth of two classes of the English population, as examples of my method of treating anthropometrical statistics. Tables I. to V. inclusive may be accepted as the highest standard of our English race, as the measurements are those of boys and men born and living under the most favourable conditions of breeding, nurture,



occupation, climate, exercise, and sanitary surroundings. They come under the first division of Class I. (persons of rank and *outdoor* professions) of the classified table of occupations given at p. 42, and are derived from the following sources: Public schools—Eton, Marlborough, Wellington, Haileybury, Clifton, Radley, and Magdalen; Britannia Training-ship for Naval Cadets; Woolwich Academy and Sandhurst College for Military Cadets; Oxford and Cambridge Universities; and St. George's Hospital Medical School.\*

Tables VI. VII. and VIII. may be accepted as a standard of the physique of the English labouring classes living in large towns, being the measurements of artisans and their children living under unfavourable sanitary conditions and occupations (Class VII. of the classified table of occupations).†

\* I am indebted to the following gentlemen for assistance in collecting these statistics: Mr. George Buck, F.R.S.; Mr. Francis Galton, F.R.S., and the members of the Anthropological Institute for reference to manuscript tables in their library; Dr. Fergus of Marlborough College; Surgeon-Major Roberts of Woolwich Academy, and Dr. Fraser of Sandhurst College; Mr. G. R. Turner of St. George's Hospital; Mr. Maclaren of Oxford; Mr. Passingham of Cambridge; and numerous private friends.

† I am indebted to the following gentlemen for assistance in collecting the materials of which these tables are chiefly composed: Mr. G. C. Steet, Telegraphic Department of the General Post Office; Dr. Crosse, Royal Military Asylum, Chelsea; Dr. F. H.

On examination it will be found that each table consists of three portions, which may be described as *index-columns*, *age-columns*, and *result-columns*. The index-columns are situated on the left and on the right of the tables, and indicate the values observed; that is to say, the heights and circumferences in inches and mètres, and the weights in pounds English and kilogrammes. The age-columns (or time- or date-columns) occupy the centre of the tables, and contain the whole number of observations for each age arranged in uniform groups. The result-columns, situated at the bottom of the tables and continuous with the age-columns, show the total number of observations, the *average* and the *mean* values in inches or pounds, and the variations or rate of growth or increase from year to year.

The total height being the most characteristic and important measurement of the body, the arrangement of the table of heights has been made the model for all the rest. As the height is taken from the ground and increases with the age of the individual till full growth is attained, the table is constructed to read from below upwards, the smaller values being placed at the bottom, and the larger

---

Baxter, Royal Hibernian School, Dublin; and Dr. Bridges, Inspector of the Local Government Board, for reference to manuscript returns made by myself and others; and to several private friends.

ones at the top—an order the reverse of that usually adopted in statistical tables referring to other subjects. In this way the table of heights follows the natural order of the development of the body from birth to maturity, and serves as a diagram as well as a record of its development. In order to obtain an adequate idea of the variations in the size of different individuals of the same age, and the limits above and below the average to which those variations extend, the whole number of observations is given;\* but as the number of observations for each age is not the same, the whole of them are reduced to a uniform rate *per thousand*, for the purpose of comparing one year with another, and further to increase the diagrammatic effect of the table. The averages are worked out from the original measurements, and not from the *per mille* groups. This is especially necessary when the materials are much condensed, as in Table II., where the weights are given in groups of 7 lbs. each.†

\* ‘Averages are numerical expressions of probabilities; extreme values are expressions of possibilities.’ Dr. Guy, in *Cyclopædia of Anatomy and Physiology*, art. ‘Statistics.’

† Weights are better arranged in groups of two or four pounds. I was obliged to adopt the groups of seven pounds because some of the statistics of the public-school boys were returned in that form, and I was unwilling to throw them aside on that account. If the weights are taken and recorded in pounds, the statistician can adopt any grouping he may find most convenient.

In Table III. the number of observations of the chest-girth are too few to be reduced to a *per mille* rate, and the original numbers are therefore given.

*Averages and Means.* It is necessary here to call attention to the difference in the meaning of the terms average and mean—which in common language are synonymous—when used in connection with anthropological inquiries (see p. 20). An *average* is obtained by dividing the sum of the values observed by the number of observations, while a *mean* is the value at which the largest number of observations occur. Thus, the average height of a number of men of different stature is obtained by multiplying the various heights by the number of men at each height and dividing the sum of their heights by the total number of men. The mean height on the other hand is obtained by arranging the men in groups, and noting the height of the group which contains the largest number of individuals. The average height is the measurement which each and all the men would possess if the tall ones could be made shorter and the short ones taller, and the predominance of a few very tall or very short men would determine the relative value of the average height. The mean height is the central or typical height which the men possess, and is the height which all of them ought to be, and probably would be, if their growth had not been inter-

fered with by accidental causes. An average includes, and is influenced by, exceptional cases, while a mean excludes and disregards exceptional cases, and is consequently uninfluenced by them. The distinction has an important practical bearing on the study of anthropology. If we wish to distinguish the specific characteristics of different races of mankind, we employ the mean; if the differences between various classes of the same race, we employ the average. Tables XII. and XIII. show the variations in the height and weight of different classes of our English community as distinguished by averages.

The physical proportions of girls and women have been very imperfectly studied in this country. The only statistics of the height, weight, and chest-girth of females which I have been able to obtain are those of girls between the ages of eight and thirteen years, of the class of artisans, agricultural labourers, and factory operatives.\* I have determined, therefore, to give in Tables X. and XI. the statistics of the height and weight of nearly eleven thousand girls between the ages of five and nineteen years, living in the city and neighbourhood of Boston, U.S. America, and published by Professor Bowditch, of Harvard University, in his 'Report on the Growth of Children.'

\* These statistics are published in my pamphlet on the *Physical Requirements of Factory Children* (Churchill, 1876).

The tables include all classes of the community, and, therefore, do not correspond with either of the classes of males given in my other tables, but they may be accepted as the mean between the two. The state of society and the school system in America does not allow of the distinction between class and class which is possible in this country. It is probably in our lower middle-class schools (Class II. of the table of occupations) that we must look for the corresponding class of males in this country. I hope that the example of collecting statistics of the physical proportions of females as well as males, adopted in America, will be followed in this country, as it is a matter of great interest and importance.

TABLE I. *Showing the actual, average, and mean HEIGHT, and the of 10 and 30 Years, of the most favoured Classes of the Cadets, Medical and University Students.*

Age last birthday.	10	11	12	13	14	15	16	17
No. of observations.	74	150	248	473	477	541	686	1602
HEIGHT (without shoes).								
From ft. in.								
From inches.								
6 5 77 to 78	—	—	—	—	—	—	—	—
6 4 76 " 77	—	—	—	—	—	—	—	1
6 3 75 " 76	—	—	—	—	—	—	—	2
6 2 74 " 75	—	—	—	—	—	—	2	2
6 1 73 " 74	—	—	—	—	—	—	2	5
6 0 72 " 73	—	—	—	—	—	—	17	25
5 11 71 " 72	—	—	—	—	—	5	18	72
5 10 70 " 71	—	—	—	—	—	17	45	98
5 9 69 " 70	—	—	—	—	8	22	79	124
5 8 68 " 69	—	—	—	—	12	28	126	158
5 7 67 " 68	—	—	—	2	14	62	156	150
5 6 66 " 67	—	—	—	6	41	72	125	139
5 5 65 " 66	—	—	—	12	58	96	139	95
5 4 64 " 65	—	—	—	14	52	122	115	64
5 3 63 " 64	—	—	8	33	83	153	63	42
5 2 62 " 63	—	—	20	42	100	116	41	14
5 1 61 " 62	—	—	21	78	129	103	30	5
5 0 60 " 61	—	13	84	130	125	76	16	2
4 11 59 " 60	—	13	84	141	136	43	13	1
4 10 58 " 59	—	60	104	151	81	34	6	—
4 9 57 " 58	13	76	133	127	67	22	6	—
4 8 56 " 57	54	140	181	97	29	19	1	1
4 7 55 " 56	122	186	177	71	27	4	—	—
4 6 54 " 55	189	220	84	61	18	3	—	—
4 5 53 " 54	230	100	52	22	8	2	—	—
4 4 52 " 53	189	93	32	8	8	1	—	—
4 3 51 " 52	122	40	7	2	4	—	—	—
4 2 50 " 51	40	46	6	2	—	—	—	—
4 1 49 " 50	14	13	4	—	—	—	—	—
4 0 48 " 49	27	—	4	—	—	—	—	—
Total.	1000	1000	1000	1000	1000	1000	1000	1000
Average height.	53.40	54.91	56.97	58.79	61.11	63.47	66.40	67.86
Average growth.	—	1.51	2.06	1.82	2.32	2.36	2.93	1.46
Mean height.	53.00	54.50	56.50	58.50	61.00	63.50	66.50	68.00
Mean growth.	—	1.50	2.00	2.00	2.50	2.50	3.00	1.50
Age last birthday.	10	11	12	13	14	15	16	17

*Annual Rate of Growth, of 7709 Boys and Men between the Ages  
English Population—Public-School Boys, Naval and Military*

18	19	20	21	22	23	24	25-30	Age last birthday.	
1522	794	391	340	205	91	45	70	No. of observations.	
								HEIGHT (without shoes).	
								Inches.	Average mètres.
1	1	—	—	—	—	—	—	77·5	1·969
1	—	2	—	—	—	—	—	76·5	1·944
2	3	7	—	4	—	—	—	75·5	1·918
4	10	10	12	6	11	—	14	74·5	1·893
19	25	25	33	10	33	22	14	73·5	1·868
34	62	79	50	53	44	44	43	72·5	1·841
73	68	94	89	78	77	111	114	71·5	1·816
124	116	135	165	146	142	133	214	70·5	1·791
136	186	184	133	137	132	134	144	69·5	1·765
167	164	148	148	157	133	267	143	68·5	1·740
142	124	115	142	142	121	133	143	67·5	1·714
122	109	86	133	146	120	67	86	66·5	1·689
69	62	61	59	49	100	45	43	65·5	1·664
54	38	38	24	35	87	44	28	64·5	1·638
39	20	10	9	34	—	—	14	63·5	1·613
7	6	—	3	4	—	—	—	62·5	1·587
4	5	—	—	—	—	—	—	61·5	1·562
1	1	—	—	—	—	—	—	60·5	1·537
1	—	—	—	—	—	—	—	59·5	1·511
—	—	—	—	—	—	—	—	58·5	1·486
—	—	—	—	—	—	—	—	57·5	1·460
—	—	—	—	—	—	—	—	56·5	1·435
—	—	—	—	—	—	—	—	55·5	1·410
—	—	—	—	—	—	—	—	54·5	1·384
—	—	—	—	—	—	—	—	53·5	1·359
—	—	—	—	—	—	—	—	52·5	1·333
—	—	—	—	—	—	—	—	51·5	1·308
—	—	—	—	—	—	—	—	50·5	1·283
—	—	—	—	—	—	—	—	49·5	1·257
—	—	—	—	—	—	—	—	48·5	1·232
1000	1000	1000	1000	1000	1000	1000	1000	Total.	
68·29	68·72	69·13	69·16	68·93	68·52	68·95	69·06	Average height.	
0·43	0·43	0·41	0·03	—	—	—	—	Average growth.	
68·50	68·75	69·00	69·00	69·00	69·00	69·00	69·00	Mean height.	
0·50	0·25	0·25	—	—	—	—	—	Mean growth.	
18	19	20	21	22	23	24	25-30	Age last birthday.	



TABLE II. *Showing the actual, average, and mean WEIGHT, and the Annual Rate of the most favoured Classes of the English Population—Public-School Boys.*

Age last birthday.		10	11	12	13	14	15	16	17
No. of observations.		74	150	248	473	477	541	696	1602
WEIGHT (including clothes = 9 lbs.).									
Stones. (1=14lbs.)	lbs. From								
14 to 15	196 to 200	—	—	—	—	—	—	—	—
13 " 14	182 " 196	—	—	—	—	—	—	—	4
12½ " 13	175 " 182	—	—	—	—	—	—	4	13
12 " 12½	168 " 175	—	—	—	—	—	—	5	27
11½ " 12	161 " 168	—	—	—	—	—	—	12	48
11 " 11½	154 " 161	—	—	—	—	—	6	44	109
10½ " 11	147 " 154	—	—	—	—	—	15	63	149
10 " 10½	140 " 147	—	—	—	—	—	23	110	170
9½ " 10	133 " 140	—	—	—	—	18	59	163	180
9 " 9½	126 " 133	—	—	—	—	12	95	174	139
8½ " 9	119 " 126	—	—	—	16	29	124	157	85
8 " 8½	112 " 119	—	—	—	17	82	122	103	45
7½ " 8	105 " 112	—	—	12	43	118	122	76	18
7 " 7½	98 " 105	—	—	36	112	247	185	44	10
6½ " 7	91 " 98	—	7	79	187	228	140	25	2
6 " 6½	84 " 91	—	67	221	256	130	61	9	1
5½ " 6	77 " 84	68	155	273	225	41	35	8	—
5 " 5½	70 " 77	270	355	233	100	38	8	3	—
4½ " 5	63 " 70	446	309	118	38	3	5	—	—
4 " 4½	56 " 63	162	100	24	6	—	—	—	—
3½ " 4	49 " 56	54	7	4	—	—	—	—	—
Total.		1000	1000	1000	1000	1000	1000	1000	1000
Average weight.		67·44	72·94	80·33	88·60	99·21	110·42	128·34	141·03
Average growth.		—	5·50	7·39	8·27	10·61	11·21	17·92	12·69
Mean weight.		67·0	73·0	80·0	88·0	98·0	110·0	126·0	140·0
Mean growth.		—	6·0	7·0	8·0	10·0	12·0	16·0	14·0
Age last birthday.		10	11	12	13	14	15	16	17

*Increase in Weight, of 7709 Boys and Men, between the Ages of 10 and 30 Years, of Naval and Military Cadets, and Medical and University Students.*

18	19	20	21	22	23	24	25-30	Age last birthday.
1522	794	391	340	205	91	45	70	No. of observations.
								WEIGHT (including clothes =9 lbs.).
								Average lbs.      Average kilos.
2	1	—	6	—	—	21	14	203·0      92·27
8	6	5	12	10	—	—	15	189·0      85·91
29	34	64	45	44	75	22	119	178·5      82·04
48	53	51	84	69	75	65	90	171·5      77·95
81	94	138	147	157	118	57	134	164·5      74·77
123	130	162	155	221	172	152	135	157·5      71·59
169	197	216	180	133	205	196	164	150·5      68·41
195	192	182	143	154	151	196	149	143·5      65·22
149	150	108	129	118	54	152	90	136·5      62·05
104	98	49	75	55	108	44	45	129·5      57·83
62	30	20	18	25	32	—	15	122·5      55·68
31	12	5	6	14	10	63	30	115·5      52·50
3	3	—	—	—	—	—	—	108·5      49·31
1	—	—	—	—	—	—	—	101·5      46·13
—	—	—	—	—	—	—	—	94·5      42·95
—	—	—	—	—	—	—	—	87·5      39·78
—	—	—	—	—	—	—	—	80·5      37·50
—	—	—	—	—	—	—	—	73·5      33·41
—	—	—	—	—	—	—	—	66·5      30·23
—	—	—	—	—	—	—	—	59·5      27·04
—	—	—	—	—	—	—	—	52·5      23·86
1000	1000	1000	1000	1000	1000	1000	1000	Total.
146·00	148·20	152·07	152·34	154·78	151·70	149·20	155·20	Average weight.
4·97	2·20	3·87	0·27	2·44	—	—	0·42	Average growth.
146·0	148·0	150·0	152·0	—	—	—	154·0	Mean weight.
6·0	2·0	2·0	2·0	—	—	—	2·0	Mean growth.
18	19	20	21	22	23	24	25-30	Age last birthday.



TABLE IV. Summary of Tables I. II. and III. Showing the average and mean Height, Weight, and Chest-girth, and the Annual Rate of Increase, from the Age of 10 to 30 Years; the Ratio of the Weight and Chest-girth to the Height (the weight or chest-girth for each inch of stature); and the Ratio of Increase (lbs. per inch and chest-girth per inch of growth in height)—Public-School Boys, Naval and Military Cadets, Medical and University Students.

Age last birth- day.	Height (without shoes).			Weight, including clothes=9 lbs.			Empty Chest-girth.			Ratio, Height=1.		Increase, Height=1.	
	Average.	Growth.	Mean.	Average.	Growth.	Mean.	Average.	Growth.	Mean.	Weight.	Chest-girth.	Weight.	Chest-girth.
	Inches.	Inches.	Inches.	Lbs.	Lbs.	Lbs.	Inches.	Inches.	Inches.	Lbs.	Inches.	Lbs.	Inches.
10	53.40	—	53.00	67.4	—	67.0	—	—	—	1.252	—	—	—
11	54.91	1.51	54.50	72.9	5.50	73.0	6.0	—	—	1.399	—	4.0	—
12	56.97	2.06	56.50	80.3	7.39	80.0	7.0	—	—	1.416	—	3.5	—
13	58.79	1.82	58.50	88.6	8.27	88.0	8.0	0.87	27.54	1.504	0.486	4.0	0.50
14	61.11	2.32	61.00	99.2	10.61	98.0	10.0	1.24	28.50	1.606	0.487	4.0	0.40
15	63.47	2.36	63.50	110.4	11.21	110.0	12.0	1.07	31.00	1.732	0.488	4.8	0.60
16	66.40	2.93	66.50	128.3	17.92	126.0	16.0	33.08	33.00	1.894	0.496	5.8	0.66
17	67.84	1.46	68.00	141.0	12.69	140.0	14.0	33.98	34.00	2.087	0.500	9.3	0.80
18	68.29	0.43	68.50	146.0	4.97	146.0	6.0	34.44	34.50	2.102	0.503	12.0	1.00
19	68.72	0.43	68.75	148.8	2.20	148.0	2.0	34.77	34.75	2.132	0.505	8.0	1.00
20	69.13	0.41	69.00	152.0	3.87	150.0	2.0	35.25	35.00	2.176	0.507	8.0	1.00
21	69.16	0.03	—	152.3	0.27	152.0	2.0	35.42	35.25	2.203	0.510	2.0	—
22	68.93	—	—	154.7	2.44	—	—	35.80	35.50	—	0.514	—	—
23	68.53	—	—	151.7	—	—	—	35.52	35.75	2.203	0.518	—	—
24	68.95	—	—	149.2	—	—	—	36.10	36.00	2.203	0.521	—	—
25-30	69.06	—	69.00	155.2	0.42	154.0	2.0	35.96	36.00	2.231	0.521	2.0	2.0

TABLE V. Table IV. reduced to *Metrical Measures and Weights (mètres are reduced to centimètres by removing the decimal point two figures to the right).*

Age last birth-day.	Height (without shoes).			Weight, including clothes=4'1.			Empty Chest-girth.			Ratio, Height=1.		Increase, Height=1.	
	Average.	Growth.	Mean.	Growth.	Average.	Mean.	Growth.	Average.	Mean.	Growth.	Weight.	Chest-girth.	Weight.
	Mètres.	Mètres.	Mètres.	Kilos.	Kilos.	Kilos.	Mètres.	Mètres.	Mètres.	Kilos.	Mètres.	—	—
10	1.357	—	1.347	—	30.64	30.45	—	—	—	22.60	—	—	—
11	1.395	0.038	1.385	2.50	33.14	33.20	2.73	—	—	23.97	—	—	—
12	1.448	0.051	1.436	3.86	36.50	36.36	3.18	—	—	25.84	—	—	—
13	1.495	0.046	1.487	3.76	40.27	40.00	3.64	0.699	0.698	26.90	0.486	—	—
14	1.553	0.059	1.550	4.82	44.54	44.54	4.54	0.722	0.724	28.73	0.487	0.005	—
15	1.613	0.060	1.614	5.09	50.18	50.00	5.45	0.753	0.750	30.98	0.488	—	—
16	1.687	0.074	1.690	8.14	58.31	57.27	7.27	0.781	0.787	33.89	0.496	—	—
17	1.724	0.037	1.728	5.77	64.09	63.64	6.36	0.840	0.838	36.93	0.500	—	—
18	1.735	0.011	1.741	0.12	66.36	66.36	2.73	0.863	0.864	38.11	0.503	—	—
19	1.747	0.011	1.748	0.006	67.40	67.27	0.90	0.875	0.876	38.48	0.505	—	—
20	1.757	0.010	1.754	0.006	69.09	68.18	0.91	0.883	0.883	38.87	0.507	—	—
21	1.758	0.003	1.754	—	69.23	69.09	0.91	0.895	0.895	39.83	0.510	—	—
22	1.752	—	1.754	—	70.82	1.11	—	0.900	0.902	—	0.514	—	—
23	1.741	—	1.754	—	69.00	—	—	0.902	0.908	—	0.518	—	—
24	1.752	—	1.754	—	68.00	—	—	0.917	0.915	—	0.521	—	—
25-30	1.754	—	1.754	—	70.54	0.19	0.91	0.913	0.915	89.31	0.521	—	—

The following statistics of the height (*length*), weight, and chest-girth of new-born infants of the artisan class are given in a separate table, as their addition to Tables VI. VII. and VIII. would make them too large for the pages on which they are printed; the measurements, moreover, are not quite identical with those of older children. The heights are taken in the recumbent position, the weights without clothes, and the chest-girths without regard to the state of the respiration. The table includes only infants born at the full period of gestation.\*

*Table showing the HEIGHT, WEIGHT, and CHEST-GIRTH of New-born Infants of the Artisan Class.*

Height.			Weight.			Chest-girth.		
Inches. From	Males.	Females	Lbs. From	Males.	Females	Inches. From	Males.	Females
23 to 24	1	1	11 to 12	1	—	15·0	1	—
22 " 23	1	—	10 " 11	—	1	14·5	1	—
21 " 22	9	4	9 " 10	6	2	14·0	4	3
20 " 21	16	9	8 " 9	31	20	13·5	4	3
19 " 20	30	25	7 " 8	29	36	13·0	6	5
18 " 19	33	50	6 " 7	27	29	12·5	1	1
17 " 18	7	10	5 " 6	4	11	12·0	2	5
16 " 17	2	1	4 " 5	1	1	11·5	—	2
15 " 16	1	—	3 " 4	1	—	11·0	1	—
—	—	—	—	—	—	10·5	—	—
—	—	—	—	—	—	10·0	—	1
Total.	100	100	Total.	100	100	Total.	20	20
Average.	19·34	18·98	Average.	7·55	7·23	Average.	13·25	12·65
Mean.	19·00	18·50	Mean.	7·50	7·00	Mean.	—	—

\* I am indebted to Dr. J. Cumming of Edinburgh and Messrs. B. W. Large and A. R. Hutchinson, House Surgeons of the Royal Maternity Hospital of that city, for these statistics. The proportions of the new-born infant necessarily form the starting-point for the study of the development of the body; and it is much to be regretted that with the many opportunities which our public institutions present, so few measurements are recorded. M. Quetelet's observations were collected at 'l'Hospice de la Maternité, sur des enfants morts, mais venus à terme et régulièrement conformés,' and as the dimensions of the dead body (still-born ?) they must differ much from those of the living model.

TABLE VI. *Showing the actual, average, and mean HEIGHT, and the Annual the Population in large Eng.*

HEIGHT (without shoes).		Age last birthday.										
		4	5	6	7	8	9	10	11	12	13	14
No. of observations.		21	175	327	781	1036	1182	1119	1080	620	991	2247
From ft. in. 6 0	From inches. 72 to 73	—	—	—	—	—	—	—	—	—	—	—
5 11	71 " 72	—	—	—	—	—	—	—	—	—	—	—
5 10	70 " 71	—	—	—	—	—	—	—	—	—	—	—
5 9	69 " 70	—	—	—	—	—	—	—	—	—	—	1
5 8	68 " 69	—	—	—	—	—	—	—	—	—	—	—
5 7	67 " 68	—	—	—	—	—	—	—	—	—	—	1
5 6	66 " 67	—	—	—	—	—	—	—	—	—	—	2
5 5	65 " 66	—	—	—	—	—	—	—	—	—	—	5
5 4	64 " 65	—	—	—	—	—	—	—	—	—	3	12
5 3	63 " 64	—	—	—	—	—	—	—	—	—	1	23
5 2	62 " 63	—	—	—	—	—	—	—	—	—	8	20
5 1	61 " 62	—	—	—	—	—	—	—	—	1	9	41
5 0	60 " 61	—	—	—	—	—	—	—	1	3	23	79
4 11	59 " 60	—	—	—	—	—	1	—	2	8	60	100
4 10	58 " 59	—	—	—	—	—	—	2	2	20	78	130
4 9	57 " 58	—	—	—	—	—	1	2	11	23	147	169
4 8	56 " 57	—	—	—	—	—	2	4	22	47	163	176
4 7	55 " 56	—	—	—	—	—	3	15	46	86	181	124
4 6	54 " 55	—	—	—	—	—	15	41	75	130	137	63
4 5	53 " 54	—	—	—	—	2	31	71	112	150	80	34
4 4	52 " 53	—	—	—	—	3	64	125	149	170	45	11
4 3	51 " 52	—	—	—	—	24	92	153	149	149	26	7
4 2	50 " 51	—	—	—	2	44	160	168	157	109	27	2
4 1	49 " 50	—	—	3	14	105	156	151	117	53	8	—
4 0	48 " 49	—	—	6	45	135	174	123	78	36	2	—
3 11	47 " 48	—	—	15	86	184	136	82	60	10	1	—
3 10	46 " 47	—	—	57	164	188	96	35	8	5	1	—
3 9	45 " 46	—	11	77	181	146	43	18	9	1	—	—
3 8	44 " 45	—	40	153	221	94	15	2	—	—	—	—
3 7	43 " 44	—	97	208	135	44	7	4	1	1	—	—
3 6	42 " 43	—	137	213	97	19	—	2	1	—	—	—
3 5	41 " 42	—	275	171	41	7	2	—	—	—	—	—
3 4	40 " 41	3	183	55	12	4	2	—	—	—	—	—
3 3	39 " 40	5	165	30	—	1	—	—	—	—	—	—
3 2	38 " 39	7	57	6	1	—	—	—	—	—	—	—
3 1	37 " 38	1	29	6	1	—	—	—	—	—	—	—
3 0	36 " 37	4	—	—	—	—	—	—	—	—	—	—
2 11	35 " 36	1	6	—	—	—	—	—	—	—	—	—
Total.		21	1000	1000	1000	1000	1000	1000	1000	1000	1000	100
Average height.		38·45	41·16	43·18	45·01	46·99	49·22	50·52	51·52	52·99	55·93	57·7
Average growth.		—	2·71	2·02	1·83	1·98	2·23	1·30	1·00	1·47	2·94	1·8
Mean height.		38·5	41·0	43·0	45·0	47·0	49·0	50·5	51·5	53·0	55·5	58·0
Mean growth.		—	2·5	2·0	2·0	2·0	2·0	1·5	1·0	1·5	2·5	2·5
Age last birthday.		4	5	6	7	8	9	10	11	12	13	14

*Rate of Growth, of 13,931 Boys and Men between the Ages of 4 and 50 Years, of Irish Towns—Artisan Class.*

Age last birthday.											HEIGHT (with- out shoes).	
15	16	17	18	19	20	21-22	23 to 30	23 to 50.				
745	1018	453	163	97	69	91	135	156	1117	318	No. of obs.	
											Average	
—	—	—	—	—	—	—	7	6	—	—	Inches.	Mètres.
—	—	—	—	—	—	—	7	6	—	—	72·5	1·841
—	—	2	6	31	14	44	7	7	—	—	71·5	1·816
1	3	6	33	21	29	33	52	45	—	—	70·5	1·791
5	11	16	33	62	44	44	97	89	—	—	69·5	1·765
8	19	40	72	82	116	77	148	160	—	—	68·5	1·740
12	39	62	80	113	174	121	126	128	—	—	67·5	1·715
14	62	106	184	144	246	308	163	161	—	—	66·5	1·689
36	88	155	139	216	116	197	156	154	—	—	65·5	1·664
55	137	182	184	155	87	99	163	160	—	—	64·5	1·638
90	154	170	157	93	73	55	22	26	—	—	63·5	1·613
95	130	122	72	42	101	22	30	32	—	—	62·5	1·587
126	117	81	20	41	—	—	29	26	—	—	61·5	1·562
118	81	15	13	—	—	—	—	6	—	—	60·5	1·537
134	63	28	6	—	—	—	—	—	—	—	59·5	1·511
89	49	11	—	—	—	—	—	—	—	—	58·5	1·486
91	33	—	—	—	—	—	—	—	—	—	57·5	1·460
66	10	4	—	—	—	—	—	—	—	—	56·5	1·435
39	3	—	2	—	—	—	—	—	—	—	55·5	1·410
13	1	—	—	—	—	—	—	—	—	—	54·5	1·384
5	—	—	—	—	—	—	—	—	—	—	53·5	1·359
3	2	—	—	—	—	—	—	—	—	—	52·5	1·333
—	—	—	—	—	—	—	—	—	—	—	51·5	1·308
—	—	—	—	—	—	—	—	—	—	—	50·5	1·283
—	—	—	—	—	—	—	—	—	—	—	49·5	1·257
—	—	—	—	—	—	—	—	—	—	—	48·5	1·232
—	—	—	—	—	—	—	—	—	Danson (Liverpool Gaol).	Dr. Beddoe (London).	47·5	1·206
—	—	—	—	—	—	—	—	—			46·5	1·181
—	—	—	—	—	—	—	—	—			45·5	1·156
—	—	—	—	—	—	—	—	—			44·5	1·130
—	—	—	—	—	—	—	—	—			43·5	1·105
—	—	—	—	—	—	—	—	—			42·5	1·079
—	—	—	—	—	—	—	—	—			41·5	1·054
—	—	—	—	—	—	—	—	—			40·5	1·028
—	—	—	—	—	—	—	—	—			39·5	1·003
—	—	—	—	—	—	—	—	—			38·5	0·978
—	—	—	—	—	—	—	—	—			37·5	0·952
—	—	—	—	—	—	—	—	—			36·5	0·927
—	—	—	—	—	—	—	—	—			35·5	0·902
1000	1000	1000	1000	1000	1000	1000	1000	—	1117	318	Total.	
60·58	62·93	64·45	65·47	66·02	66·31	66·60	66·68	66·65	66·39	66·72	Av. height.	
2·82	2·35	1·52	1·02	0·55	0·29	0·29	0·08	—	—	—	Av. growth.	
60·5	63·0	64·5	65·5	66·0	66·25	66·5	—	66·5	—	—	Mean height.	
2·5	2·5	1·5	1·0	0·5	0·25	0·25	—	—	—	—	„ growth.	
15	16	17	18	19	20	21-22	23-30	23 to 50			Age 1. birth.	



TABLE VII. *Showing the actual, average, and mean WEIGHT, and the Annual the Population in large*

WEIGHT (including clothes).		Age last birthday.									
		4	5	6	7	8	9	10	11	12	13
No. of observations.		21	176	327	631	1038	1203	1126	979	615	1054
Stones (14 lbs.).	Lbs. From										
13 to 14	182 to 196	—	—	—	—	—	—	—	—	—	—
12½ " 13	175 " 182	—	—	—	—	—	—	—	—	—	—
12 " 12½	168 " 175	—	—	—	—	—	—	—	—	—	—
11½ " 12	161 " 168	—	—	—	—	—	—	—	—	—	—
11 " 11½	154 " 161	—	—	—	—	—	—	—	—	—	—
10½ " 11	147 " 154	—	—	—	—	—	—	—	—	—	—
10 " 10½	140 " 147	—	—	—	—	—	—	—	—	—	—
9½ " 10	133 " 140	—	—	—	—	—	—	—	—	—	—
9 " 9½	126 " 133	—	—	—	—	—	—	—	—	—	—
8½ " 9	119 " 126	—	—	—	—	—	—	—	—	—	—
8 " 8½	112 " 119	—	—	—	—	—	—	—	—	—	3
7½ " 8	105 " 112	—	—	—	—	—	—	—	—	—	4
7 " 7½	98 " 105	—	—	—	—	—	—	1	5	20	—
6½ " 7	91 " 98	—	—	—	—	—	1	6	31	52	—
6 " 6½	84 " 91	—	—	—	—	1	12	41	84	171	—
5½ " 6	77 " 84	—	—	—	—	23	64	135	267	280	—
5 " 5½	70 " 77	—	—	6	32	106	206	304	312	304	—
4½ " 5	63 " 70	—	6	24	144	216	253	386	262	228	120
4 " 4½	56 " 63	—	108	355	419	451	335	268	201	68	40
3½ " 4	49 " 56	1	534	477	332	236	160	59	48	5	6
3 " 3½	42 " 49	9	239	126	97	62	22	4	2	—	—
2½ " 3	35 " 42	8	102	18	2	2	—	—	—	—	—
2 " 2½	28 " 35	3	11	—	—	—	—	—	—	—	—
Total.		21	1000	1000	1000	1000	1000	1000	1000	1000	1000
Average weight.		41·16	49·99	54·19	56·89	59·00	62·56	66·31	69·46	73·68	78·27
Average increase.		—	8·83	4·20	2·70	2·11	3·56	3·75	3·15	4·22	4·59
Mean weight.		44·0	50·0	54·0	57·0	59·0	62·0	66·0	70·0	74·0	78·0
Mean increase.		—	6·0	4·0	3·0	2·0	3·0	4·0	4·0	4·0	4·0
Age last birthday.		4	5	6	7	8	9	10	11	12	13

*Rate of Increase of Boys and Men, between the Ages of 4 and 50 Years, of English Towns—Artisan Class.*

Age last birthday.										WEIGHT (including clothes).	
14	15	16	17	18	19	20	21-22	23 to 30	23 to 50	No. of observations.	
2094	910	1038	504	147	105	63	93	121	142	Average lbs.	Average kilos.
—	—	—	—	—	—	—	—	8	7	189·5	85·91
—	—	—	—	6	9	—	—	8	14	178·5	82·04
—	—	—	—	—	—	—	32	33	35	171·5	77·95
—	—	—	—	13	—	—	32	42	49	164·5	74·77
—	—	1	6	15	19	15	65	99	106	157·5	71·59
—	—	1	10	27	57	73	32	58	71	150·5	68·41
—	5	8	30	48	124	132	183	198	190	143·5	65·22
—	6	36	47	75	153	177	161	182	162	136·5	62·05
2	19	77	135	251	200	294	269	165	162	129·5	57·83
4	35	123	159	170	124	132	86	116	106	122·5	55·68
15	85	175	278	224	219	118	118	75	70	115·5	52·50
34	123	160	137	75	76	44	22	8	14	108·5	49·31
59	155	185	125	61	19	15	—	8	7	101·5	46·13
106	188	107	41	14	—	—	—	—	7	94·5	42·95
250	204	101	24	8	—	—	—	—	—	87·5	39·78
288	113	19	6	7	—	—	—	—	—	80·5	37·50
186	58	6	2	6	—	—	—	—	—	73·5	33·41
48	9	1	—	—	—	—	—	—	—	66·5	30·23
6	—	—	—	—	—	—	—	—	—	59·5	27·04
2	—	—	—	—	—	—	—	—	—	52·5	23·86
—	—	—	—	—	—	—	—	—	—	45·5	20·68
—	—	—	—	—	—	—	—	—	—	38·5	17·50
—	—	—	—	—	—	—	—	—	—	31·5	14·32
1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	Total.	
84·61	96·79	108·7	116·4	123·3	128·4	130·6	135·4	139·0	141·2	Average weight.	
6·34	12·18	11·93	7·66	6·97	5·08	2·20	4·81	3·58	5·74	Average increase.	
84·0	94·0	106·0	116·0	122·0	128·0	132·0	136·0	138·0	140·0	Mean weight.	
6·0	10·0	12·0	10·0	6·0	6·0	4·0	4·0	2·0	4·0	Mean increase.	
14	15	16	17	18	19	20	21-22	23-30	23-50	Age last birthday.	

TABLE VIII. *Showing the CHEST-GIRTH*

Empty CHEST-GIRTH.		Age last birthday.							
		5	6	7	8	9	10	11	12
No. of observations.		128	287	728	874	824	783	840	426
Inches. From	Average inches.								
37 to 38	37.5	—	—	—	—	—	—	—	—
36 " 37	36.5	—	—	—	—	—	—	—	—
35 " 36	35.5	—	—	—	—	—	—	—	—
34 " 35	34.5	—	—	—	—	—	—	—	—
33 " 34	33.5	—	—	—	—	—	—	—	—
32 " 33	32.5	—	—	—	—	—	—	—	—
31 " 32	31.5	—	—	—	—	—	—	—	—
30 " 31	30.5	—	—	—	—	—	—	—	1
29 " 30	29.5	—	—	—	—	—	1	2	5
28 " 29	28.5	—	—	—	—	1	12	2	11
27 " 28	27.5	—	—	—	5	24	42	21	64
26 " 27	26.5	—	—	3	40	75	108	80	171
25 " 26	25.5	—	22	54	129	177	195	207	303
24 " 25	24.5	10	80	177	269	242	258	312	303
23 " 24	23.5	31	80	242	301	268	211	242	121
22 " 23	22.5	227	258	332	258	211	203	126	111
21 " 22	21.5	391	415	236	176	73	66	13	9
20 " 21	20.5	320	213	82	30	5	1	3	2
19 " 20	19.5	31	24	6	1	—	—	—	—
Total.		1000	1000	1000	1000	1000	1000	1000	1000
Average Chest-girth.		21.40	21.68	22.54	23.09	23.79	24.08	24.34	25.24
Average increase.		—	0.28	0.86	0.55	0.70	0.29	0.46	0.31
Mean Chest-girth.		21.0	21.5	22.0	22.5	23.0	23.5	24.0	25.0
Mean increase.		—	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Age last birthday.		5	6	7	8	9	10	11	12

*of English Town Population—Artisan Class.*

Age last birthday.										Empty CHEST-GIRTH.
14	15	16	17	18	19	20	21 to 22	23 to 30	23 to 50	
1133	514	643	376	168	90	46	45	88	106	No. of observations.
—	—	—	—	—	—	—	—	—	10	Average metres.
—	—	—	—	—	—	—	—	23	19	0·952
—	—	—	—	—	11	—	22	45	48	0·927
—	—	9	—	6	—	21	67	91	114	0·902
—	2	12	2	30	12	22	111	170	171	0·876
1	10	23	40	77	178	152	178	273	257	0·851
2	19	67	83	131	211	261	222	193	172	0·826
7	76	165	213	267	198	348	245	148	133	0·800
36	84	208	309	261	222	109	133	23	38	0·775
92	185	212	167	156	145	44	23	34	38	0·749
183	193	170	131	42	12	22	—	—	—	0·724
246	243	98	32	24	11	21	—	—	—	0·698
223	124	31	21	6	—	—	—	—	—	0·673
136	50	5	2	—	—	—	—	—	—	0·648
64	14	—	—	—	—	—	—	—	—	0·623
9	—	—	—	—	—	—	—	—	—	0·617
1	—	—	—	—	—	—	—	—	—	0·571
—	—	—	—	—	—	—	—	—	—	0·546
—	—	—	—	—	—	—	—	—	—	0·521
—	—	—	—	—	—	—	—	—	—	0·495
1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	Total.
26·28	27·51	28·97	29·38	30·07	30·56	30·86	31·61	32·38	32·62	Av. Chest-girth.
1·04	1·23	1·46	0·41	0·69	0·49	0·30	0·75	0·77	0·24	Average increase.
26·0	27·0	28·5	29·5	30·0	30·5	31·0	31·5	32·0	32·5	Mean Chest-girth.
1·0	1·0	1·5	1·0	0·5	0·5	0·5	0·5	0·5	0·5	Mean increase.
14	15	16	17	18	19	20	21-22	23-30	23-50	Age last birthday.

TABLE IX. Summary of Tables VI. VII. and VIII. Showing the Height, Weight, and Chest-girth of English Town Population—Artisan Class.

Age last birth-day.	Height without shoes.				Weight incl. clothes=7 and 10 lbs.				Empty Chest-grth.				Ratio, Height=1.		Increase, Height=1.				
	Average, inches.	Growth, inches.	Mean, inches.	Growth, inches.	Average, lbs.	Growth, lbs.	Mean, lbs.	Growth, lbs.	Average, inches.	Growth, inches.	Mean, inches.	Growth, inches.	Weight.	Chest-grth.					
Birth.	19.34	19.50	19.00	19.50	7.55	33.61	7.50	36.0	13.25	8.15	—	—	0.395	0.697	—				
—	—		—		—	—	—		—		—	—	—	—	—	—	—	—	
4	38.45		38.80		—	41.16	8.83		44.0		6.0	21.40	21.0	21.0	0.5	1.143	—	—	—
5	41.16		2.71		41.00	2.50	49.99		50.0		50.0	21.68	21.5	21.5	0.5	1.220	0.512	2.40	0.25
6	43.18	2.02	43.00	2.00	54.19	54.0	54.0	4.0	21.68	0.28	21.5	0.5	1.255	0.500	2.00	0.25			
7	45.01	1.83	45.00	2.00	56.89	57.0	57.0	3.0	22.54	0.86	22.0	0.5	1.266	0.488	1.50	0.25			
8	46.99	1.98	47.00	2.00	59.00	59.0	59.0	2.0	23.09	0.55	22.5	0.5	1.265	0.482	1.00	0.25			
9	49.22	2.23	49.00	2.00	62.56	62.0	62.0	3.0	23.79	0.70	23.0	0.5	1.265	0.469	1.50	0.25			
10	50.52	1.30	50.50	1.50	66.31	66.0	66.0	4.0	24.08	0.29	23.5	0.5	1.307	0.402	2.66	0.33			
11	51.52	1.00	51.50	1.00	69.46	69.0	69.0	4.0	24.34	0.46	24.0	0.5	1.359	0.466	4.00	0.50			
12	52.99	1.47	53.50	1.50	73.68	74.0	74.0	4.0	24.93	0.89	24.5	0.5	1.383	0.460	2.66	0.33			
13	55.93	2.94	55.50	2.50	78.27	78.0	78.0	4.0	26.24	1.01	25.0	0.5	1.405	0.450	1.60	0.20			
14	57.76	1.83	58.00	2.50	84.61	84.0	84.0	6.0	26.28	0.34	26.0	1.0	1.448	0.448	2.40	0.40			
15	60.58	2.82	60.50	2.50	96.79	96.0	96.0	10.0	27.51	1.23	27.0	1.0	1.553	0.446	4.00	0.40			
16	62.93	2.35	63.00	2.50	108.7	108.0	108.0	12.0	28.97	1.46	28.5	1.5	1.682	0.452	4.80	0.60			
17	64.45	1.52	64.50	1.50	116.4	116.0	116.0	10.0	29.38	0.41	29.5	1.0	1.798	0.452	6.66	0.66			
18	65.47	1.02	65.50	1.00	128.3	128.0	128.0	6.0	30.07	0.69	30.0	0.5	1.879	0.535	6.00	0.50			
19	66.02	0.55	66.00	0.50	128.4	128.0	128.0	6.0	30.56	0.49	30.5	0.5	1.940	0.460	12.00	1.00			
20	66.31	0.29	66.25	0.25	130.6	130.0	130.0	4.0	30.87	0.80	31.0	0.5	1.992	0.467	16.00	2.00			
21	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
22	66.60	0.29	66.50	0.25	135.4	135.0	135.0	4.0	30.61	0.75	31.5	0.5	2.045	0.473	16.00	2.00			
23.30	66.68	0.08	66.50	—	139.0	138.0	138.0	2.0	32.38	0.77	32.0	0.5	2.075	0.481	2.00	—			
23.50	66.65	—	66.50	—	141.2	140.0	140.0	4.0	32.62	0.24	32.5	0.5	2.105	0.490	4.00	1.00			
Danson.	66.39	—	—	—	143.7	—	—	—	—	—	—	—	—	—	—	—			
Beddoe.	66.72	0.07	—	—	137.7	—	—	—	—	—	—	—	—	—	—	—			

The following tables of the heights and weights of girls living in Boston, U.S.A., and neighbourhood, include 3681 of American, 3623 of Irish, 585 of German, and 1397 of mixed English, Irish, and American parentage. The results of Dr. Bowditch's analysis of the racial elements is, that the children of American-born parents (and this is true of both sexes) are taller and heavier than those of foreign origin. This superiority in size he attributes partly to difference of race and climate, and partly to the greater comfort and better nurture of the older residents of Boston compared with that of the poorer immigrants from Europe. He makes no note of the Scotch element, which must be largely represented in Boston, as it is in most American towns, whether in Canada or the United States, and which in both height and weight exceeds both the English and the Irish at home.\*

\* Professor Bowditch has, at my suggestion, tabulated his statistics in the manner I am now advocating (a plan which I first adopted in my paper on Factory Children in the *Jour. Statist. Soc.* 1876); hence the facility with which they can be compared with my own. His printer has, however, omitted a portion of the *index-columns* as unnecessary, and supplied the place of the absent figures by dashes. The consequence is that the two indexes do not agree, and the children gain or lose an inch in height and four pounds in weight according to the use we make of the right or left hand columns in reading the tables. The indexes on the left are obviously the correct ones, and I have adopted them in the following tables; but the error is one which might lead to much confusion, and Dr. Bowditch will be glad to have it pointed out, as I am also glad to quote it to confirm what I have already insisted on, namely, the absolute necessity of making statistical tables complete in every detail, so that no doubt or mistake can arise in reading them.

TABLE X. *Showing the actual, average, and mean HEIGHT, America, between the Ages of 5 and 19 Years. 'The of the City Proper (Boston, U.S.A.), in several Schools in Miss Hubbard's School for Young Ladies, and in of Children,' by H. P. Bowditch, M.D.; Eighth Annual*

HEIGHT (without shoes).		Age last birthday.						
		5	6	7	8	9	10	11
No. of observations.		605	987	1139	1299	1149	1089	936
From ft. in.	From inches.							
5 9	69 to 70	—	—	—	—	—	—	—
5 8	68 " 69	—	—	—	—	—	—	—
5 7	67 " 68	—	—	—	—	—	—	—
5 6	66 " 67	—	—	—	—	—	—	—
5 5	65 " 66	—	—	—	—	—	—	—
5 4	64 " 65	—	—	—	—	—	—	—
5 3	63 " 64	—	—	—	—	—	—	1
5 2	62 " 63	—	—	—	—	—	—	2
5 1	61 " 62	—	—	—	—	—	—	1
5 0	60 " 61	—	—	—	—	—	—	3
4 11	59 " 60	—	—	—	—	—	1	15
4 10	58 " 59	—	—	—	1	—	4	26
4 9	57 " 58	—	—	—	1	1	9	54
4 8	56 " 57	—	—	—	—	2	24	71
4 7	55 " 56	—	—	—	1	7	34	112
4 6	54 " 55	—	—	—	—	17	65	137
4 5	53 " 54	—	—	—	6	26	105	133
4 4	52 " 53	—	—	2	14	69	163	169
4 3	51 " 52	—	—	3	42	114	174	117
4 2	50 " 51	—	3	10	78	170	163	81
4 1	49 " 50	—	3	36	130	186	113	42
4 0	48 " 49	—	6	56	165	166	70	24
3 11	47 " 48	1	22	147	179	98	48	8
3 10	46 " 47	7	54	170	176	78	14	2
3 9	45 " 46	28	116	204	101	35	9	—
3 8	44 " 45	33	163	167	60	12	1	—
3 7	43 " 44	119	224	119	30	4	1	1
3 6	42 " 43	190	199	61	11	2	1	—
3 5	41 " 42	212	122	16	3	2	1	1
3 4	40 " 41	198	55	6	1	1	—	—
3 3	39 " 40	121	22	2	—	1	—	—
3 2	38 " 39	56	8	1	—	—	—	—
3 1	37 " 38	32	2	—	1	—	—	—
3 0	36 " 37	3	1	—	—	—	—	—
Total.		1000	1000	1000	1000	1000	1000	1000
Average height.		41·19	43·35	45·52	47·58	49·39	51·34	53·42
Average growth.		—	2·16	2·17	2·06	1·79	1·97	2·08
Mean height.		41·0	43·5	45·5	47·5	49·5	51·5	53·5
Mean growth.		—	2·5	2·0	2·0	2·0	2·0	2·0
Age last birthday.		5	6	7	8	9	10	11

*and the Annual Rate of Growth, of 10,904 GIRLS in U.S. Statistical Data were collected in nearly all the Public-Schools in South Boston, Roxbury, Charleston, and Jamaica Plain, several of the Public Schools of Brooklyn.' See the 'Growth Report State Board of Health of Mass., Boston, U.S.A., 1877.*

Age last birthday.							HEIGHT (without shoes).	
12	13	14	15	16	17	18		
935	830	675	459	353	233	155	10,904	
							Average	
							Inches.	Mètres.
—	1	—	—	—	4	—	69·5	1·765
—	—	1	—	—	4	6	68·5	1·740
1	1	2	6	14	13	26	67·5	1·715
1	—	4	13	14	22	13	66·5	1·689
2	4	11	31	54	47	39	65·5	1·664
3	7	34	59	76	82	135	64·5	1·638
6	29	47	107	133	163	71	63·5	1·613
10	38	92	124	173	202	181	62·5	1·587
19	90	153	203	161	150	194	61·5	1·562
39	117	196	174	145	107	155	60·5	1·537
70	139	148	124	105	116	116	59·5	1·511
81	122	126	83	68	52	45	58·5	1·486
109	132	86	44	37	30	13	57·5	1·460
142	98	47	15	17	4	6	56·5	1·435
137	87	27	13	—	4	—	55·5	1·410
119	70	16	4	—	—	—	54·5	1·384
111	39	5	—	—	—	—	53·5	1·359
65	13	4	—	—	—	—	52·5	1·333
46	7	1	—	—	—	—	51·5	1·308
23	3	—	—	—	—	—	50·5	1·283
8	1	—	—	—	—	—	49·5	1·257
6	1	—	—	—	—	—	48·5	1·232
2	—	—	—	—	—	—	47·5	1·206
—	1	—	—	—	—	—	46·5	1·181
—	—	—	—	—	—	—	45·5	1·156
—	—	—	—	—	—	—	44·5	1·130
—	—	—	—	—	—	—	43·5	1·105
—	—	—	—	—	—	—	42·5	1·079
—	—	—	—	—	—	—	41·5	1·054
—	—	—	—	—	—	—	40·5	1·028
—	—	—	—	—	—	—	39·5	1·003
—	—	—	—	—	—	—	38·5	0·978
—	—	—	—	—	—	—	37·5	0·952
—	—	—	—	—	—	—	36·5	0·927
1000	1000	1000	1000	1000	1000	1000	Total.	
55·88	58·16	59·94	61·10	61·59	61·92	61·95	Average height.	
2·46	2·26	1·78	1·16	0·49	0·33	0·03	Average growth.	
56·0	58·0	60·0	61·0	61·5	62·0	62·0	Mean height.	
2·5	2·0	2·0	1·0	0·5	0·5	—	Mean growth.	
12	13	14	15	16	17	18	Age last birthday.	



TABLE XI. *Showing the actual, average, and mean WEIGHT, of America, between the Ages of*

WEIGHT (including clothes).		Age last birthday.						
		5	6	7	8	9	10	11
No. of observations.		605	987	1199	1299	1149	1089	936
Stones.	From							
	Lbs.							
11·0	158 to 222	—	—	—	—	—	—	—
	154 „ 158	—	—	—	—	—	—	—
9·0	150 „ 154	—	—	—	—	—	—	—
	146 „ 150	—	—	—	—	—	—	—
	142 „ 146	—	—	—	—	—	—	—
	138 „ 142	—	—	—	—	—	—	—
	134 „ 138	—	—	—	—	—	—	—
	130 „ 134	—	—	—	—	—	—	—
	126 „ 130	—	—	—	—	—	—	—
7·0	122 „ 126	—	—	—	—	—	—	—
	118 „ 122	—	—	—	—	—	—	1
	114 „ 118	—	—	—	—	—	—	1
	110 „ 114	—	—	—	—	—	—	2
	106 „ 110	—	—	—	—	1	1	4
	102 „ 106	—	—	—	—	1	3	5
	98 „ 102	—	—	—	—	—	1	7
5·0	94 „ 98	—	—	—	—	1	2	4
	90 „ 94	—	—	—	1	—	5	19
	86 „ 90	—	—	—	—	1	10	34
	82 „ 86	—	—	—	—	4	8	39
	78 „ 82	—	—	—	—	3	18	60
	74 „ 78	—	—	—	3	7	41	102
	70 „ 74	—	1	—	6	31	76	139
3·0	66 „ 70	—	1	2	10	45	143	140
	62 „ 66	—	1	9	36	123	159	160
	58 „ 62	—	3	28	106	197	218	139
	54 „ 58	5	16	91	203	248	168	96
	50 „ 54	21	68	178	263	194	104	35
	46 „ 50	45	180	284	219	98	34	13
	42 „ 46	201	317	265	122	39	6	—
	38 „ 42	384	287	125	26	6	2	—
	34 „ 38	265	118	17	5	1	1	—
	30 „ 34	76	8	1	—	—	—	—
	26 „ 30	3	—	—	—	—	—	—
Total.		1000	1000	1000	1000	1000	1000	1000
Average weight.		39·66	43·28	47·46	52·04	57·07	62·35	68·84
Average growth.		—	3·62	4·18	4·58	5·03	5·28	6·49
Mean weight.		40·0	44·0	48·0	52·0	56·0	60·0	66·0
Mean growth.		—	4·0	4·0	4·0	4·0	4·0	6·0
Age last birthday.		5	6	7	8	9	10	11

*and the Annual Rate of Increase, of 10,904 GIRLS in U.S.  
5 and 19 Years (see Table X.).*

Age last birthday							WEIGHT (including clothes).	
12	13	14	15	16	17	18		
935	830	675	459	353	293	155	10,904	
1	1	2	2	5	17	17	Average	
							Lbs.	Kilos.
—	—	3	2	3	4	—	190	86.4
							156	70.9
—	—	1	2	5	—	—	152	69.1
—	1	3	—	3	9	—	148	67.3
—	—	—	8	3	13	26	144	65.4
—	—	6	7	11	34	6	140	63.6
1	1	7	19	37	43	39	136	61.8
1	1	3	20	42	43	71	132	60.0
3	6	18	22	68	61	58	128	58.2
3	6	18	37	71	95	103	124	56.4
3	16	27	74	99	69	58	120	54.5
6	29	50	68	93	70	65	116	52.7
11	24	59	96	85	142	161	112	50.9
6	41	95	109	108	163	65	108	49.0
18	77	85	153	105	86	142	104	47.3
19	65	116	136	82	48	97	100	45.4
33	84	107	72	74	47	58	96	43.6
57	95	104	63	40	30	13	92	41.8
85	107	92	37	40	5	13	88	40.0
108	88	74	37	9	17	6	84	38.2
120	87	74	15	11	—	—	80	36.4
109	100	18	15	3	4	—	76	34.5
138	78	22	6	3	—	—	72	32.7
117	41	10	—	—	—	—	68	30.9
81	29	5	—	—	—	—	64	29.1
53	17	1	—	—	—	—	60	27.3
21	5	—	—	—	—	—	56	25.4
5	1	—	—	—	—	—	52	23.6
1	—	—	—	—	—	—	48	21.8
—	—	—	—	—	—	—	44	20.0
—	—	—	—	—	—	—	40	18.2
—	—	—	—	—	—	—	36	16.4
—	—	—	—	—	—	—	32	14.5
—	—	—	—	—	—	—	28	12.7
1000	1000	1000	1000	1000	1000	1000	Total.	
78.31	88.65	98.43	106.0	112.0	115.5	115.1	Average weight.	
9.47	10.34	9.78	7.65	5.95	3.50	—	Average growth.	
76.0	88.0	96.0	104.0	110.0	112.0	114.0	Mean weight.	
10.0	12.0	8.0	8.0	6.0	2.0	2.0	Mean growth.	
12	13	14	15	16	17	18	Age last birthday.	

## DIAGRAMS AND CHART-TRACINGS.

The *Chart-tracings* given at pages 96 and 112 will illustrate the manner of using the ruled portion of the Anthropometrical Chart. Diagram I. shows the mean height, weight, and chest-girth, and the relation of the weight to the height of the boys and men enumerated in Tables I. II. and III. Diagram II. shows the growth of the body in height from birth to thirty years of age; and Diagram III. its development in transverse diameter and in circumference.

The tracings require little explanation: Diagram I. reads from left to right; the names of the parts measured and the weights are, according to the directions, given in the charts and written above the tracings; and the reference numbers to the Systematic Table of Measurements are also given. The dotted line representing the weight of the body without clothes has been constructed by deducting 9 lbs. for the weight of the clothes, which were included when the observations were made.

Diagram II. reads across the page from left to right, and shows the annual rate of growth from birth to thirty years of age of the parts indicated by the names and the reference numbers in a typical or mean man in this country.

Diagram III. reads from the middle of the page to the right and left, and shows the rate of increase of

the body in breadth and circumference during the same period of life. The black lines of Diagram III. represent a transverse section of the body at birth, and at the ages of 1, 3, 7, 15, and 30 years; or they show the space which would be covered by the body if laid flat on the table, by the head, the shoulders, the chest, the abdomen, and the hips. The dotted lines show the circumferences of the same parts—*i.e.* they indicate the ends of a tape which has been passed round the body and spread out on the table. The proper distances between the parts measured are not, of course, given, as the variations of the heights with age would prevent the parallelism of the lines. It will be understood that similar tracings can be made of any other parts of the body—the arms, legs, &c.

NOTE. Artists engaged in representing the human form may accept the tracings on Diagrams II. and III. as scales of proportions of the body, not necessarily for exact imitation, but as supplying measurements which will help them to find the relative proportions of the body, and place some reasonable limits to their imaginations. Year by year many of our English artists are departing farther and farther from the natural proportions, and the formulæ adopted and taught by the most distinguished artists of the Renaissance. A passing visit to the National Gallery and to the Royal Academy and the Grosvenor Galleries will sufficiently demonstrate the truth of this assertion, and it is unnecessary to specify any particular examples.

## CHAPTER VII.

### THE PHYSICAL DEVELOPMENT\* AND THE PROPORTIONS OF THE HUMAN BODY.

THE following summary of what we already know of the development and the proportions of the body will be useful to the student of anthropometry, and indicate the nature and direction of his labours. The art-student may also consult this chapter with advantage.†

A glance at the tables will show how impossible it is to study the progressive development of man, and the causes which check or promote it, by the use of mere averages. In a thousand men or boys the average represents only a small fraction of the

\* 'The term growth ought strictly to be confined to mere increase of size, and development to changes of structure. Now, a child is said to grow into a man, and a foal into a horse; but, as in these cases there is much change of structure, the process properly belongs to the order of development' (p. 389). 'The child, strictly speaking, does not grow into the man, but includes germs which slowly and successively become developed and form the man' (p. 404). *The Variation of Animals and Plants under Domestication*. By C. Darwin, M.A., F.R.S. (London, 1868).

† For further details, see Quetelet's *Anthropometry*, page 253.

whole number of the same age, and differs from groups above and below it by containing only a few more individuals. The difference in height between an average public-school boy of the age of  $13\frac{1}{2}$  and  $14\frac{1}{2}$  is about 2 inches, but the difference between the tallest and shortest boys of either of these ages is 20 inches, while at birth they differed to the extent of not more than 9 inches, and probably much less. As I have already observed, it is the variations of the individuals above and below the average or mean which is significant of the agencies at work in modifying the development of the body; and it is for this reason that I have given the whole range of heights which I have found to occur at each age, and which explains the formidable array of figures which the tables present.

The permanent and constant elements which modify the development of the human body are age, sex, and race, and some of the secondary and temporary ones are disease, occupation, social habits, nurture, food, exercise, rest, &c.

In my tables no attempt has been made to distinguish the original racial elements; but Dr. J. Beddoe has shown on the adult population that the Northern or Scandinavian element is larger than the Teutonic, and the Teutonic than the Celtic.

The tables sufficiently indicate the changes which

take place in the height, according to the age; and the double series of observations for each age of the artisan and non-labouring classes, show the modifying influences of social surroundings on the development. During the periods over which the observations extend, the most favoured class has a mean height of about two inches greater than the industrial class, but it is probable that if the observations were extended to the time of birth they would gradually approximate, and become identical at that epoch.


The total height of the new-born infant differs but little in the two sexes, the average for boys being 19·34 inches, and for girls 18·98 inches, the difference being rather less than half an inch. This difference, slight in infancy, maintains itself till near the age of thirteen years, when, in this country and America at least, the average girl is taller and heavier than the boy. This halting, so to speak, of the male, is speedily recovered; and the superiority is maintained to the period of full development, when the proportions of the two sexes stand as 1 to 0·937, or of 16 to 15.

Quetelet's observations show that both men and women maintain their maximum height till the age of fifty years, when it begins to recede, and at the age of ninety has lost about  $1\frac{1}{2}$  inches. This diminution of the height by advancing years is to be at






I. Diagram showing tracings on the Chart of  
in Tables I, II, and III, and the



The mean Heights, Chest - girths, and Weights, given  
 the relation of the Weight to the Height.


between the birthdays, i.e.  $10\frac{1}{2}$ ,  $11\frac{1}{2}$ ,  $12\frac{1}{2}$  &c. )








## II. Diagram showing the Deve



*elop*





tributed to the contraction of the soft parts between the ends of the bones, especially the cartilages of the spinal column, and to the stooping position assumed by old people. In considering the absolute height, the growth is most rapid in the first year, when it amounts to about  $7\frac{3}{4}$  inches; during the second year it is already reduced one-half, being a little less than 4 inches; and it becomes gradually less till the age of twelve, when it is reduced to about  $1\frac{1}{2}$  or 2 inches, according to the social condition. With the accession of puberty there is an increased rate of growth in the non-labouring classes, and an entire cessation of it at the age of nineteen or twenty years; while in the artisan classes the growth is more uniform, and extends to about the twenty-third year.

The height of woman is less than that of man, for three different causes: firstly, the woman is born a little smaller; secondly, her annual rate of growth after the age of fourteen is feebler; and thirdly, her growth terminates about two years before that of man. It is to this last cause above all that the difference in height must be attributed; for while at the age of thirteen girls are a little taller than boys, at the period of full development women are nearly 4 inches shorter than men.

‘In considering a particular individual,’ says M. Quetelet, ‘his growth is far from being as regular as



that indicated in the tables. There are nearly always points at which the development of a person is arrested, as also there are times of growth more or less rapid. These anomalies are observed about the age of puberty and after illness. Favourable circumstances would have to be simultaneous for all the physical faculties to develop themselves in a perfectly regular manner. When we look on a great number of persons these little anomalies disappear in the general means, and what is wanting in the development of one is compensated for by an excess of growth in another.'

About the age of fifteen a rapid development in height takes place. This growth precedes puberty, and a similar change shows itself in the girls, but from one to two years earlier. It is probably to the greater or less development of the body at the time of the accession of puberty that the final difference in the height of individuals is chiefly to be attributed; hence the influences which promote or retard growth at this period are most deserving of study. In boys puberty occurs later, and is less regular and decided, than in girls. The transition from boyhood to manhood extends over a period of three or four years, and is accompanied by increased physical development of the body; but girls develop into women in a few months, and with the complete establishment of puberty growth in height is much diminished, and

often ceases altogether. This subject would be best illustrated by making a series of successive annual measurements of the same individuals (especially of girls), a labour, however, which could only be effectually carried out by a large number of observers.

The tables of heights of boys show that the pre-pubertic growth begins about the age of  $12\frac{1}{2}$  or 13 years in the non-labouring class, and causes not only a higher mean, but a wider range, between the tallest and shortest boys than among the labouring class, amongst whom the pre-pubertic growth does not show itself till a year or two later.

The contrast presented by the columns of figures representing the non-labouring and the artisan class from the age of twelve to seventeen shows the marked effect of social surroundings on the development of the body; the one class being retarded and depressed by laborious occupations and insanitary influences, the other expanded, and probably exaggerated, by the prevalence of circumstances favourable to growth. The position of the body influences the stature. The upright position retards development in height, while the recumbent position—even though nutrition may be impaired by disease—favours increase of stature, as we often see in delicate children.

Tables XII. and XIII. show the average height and weight of different classes of the English popu-

lation arranged on the plan recommended at page 42. Although each table comprises upwards of 54,000 observations the numerous blank spaces, especially at ages below five years, show how much labour remains to be done before we can arrive at safe conclusions as to the relative physique of different classes of the community.

The materials forming these tables have been derived from the following sources: The first division of Class I. is given in detail in Tables I. and II., and is explained at page 65. The second division of Class I. are chiefly townbred boys attending the following town schools: Liverpool College; King Edward's School, Birmingham; City of London School; and Christ's Hospital. I have not been able to collect any statistics of Class II., but they will probably be found to differ little from the second division of Class I., just mentioned.

Class III. are the children of soldiers in the Royal Military Asylum, Chelsea, and the Royal Hibernian Asylum, Dublin; and recruits for the army in 1862-3. For these returns I am indebted to Drs. Crosse and Baxter, the Medical Superintendents of the Asylums; and for the recruits to Dr. Crosse's very intelligent Hospital Sergeant, Mr. P. Coughlan, who has also helped me in working out some of the averages of my tables.

Classes IV. and VIII. comprise the statistics collected by myself and other assistant commissioners for the Local Government Board, for the *Report on the Changes in the Hours and Ages of Employment of Children in Factories*, previous to the recent alterations in the Factory Acts.

The boys in Class V. are the children of sailors and coastguard men, being educated and trained for the navy in the Royal Hospital School, Greenwich, obtained through Mr. G. W. Armstrong, the resident medical officer.

The statistics in Class VII. are derived from various sources in large towns. Many of the boys were the applicants for appointments as messengers and clerks in the Telegraph Department of the General Post Office, and the rejected as well as the accepted candidates are included. I am indebted to Mr. Steet, the medical officer, for these returns.

The averages at the ages from twenty-three to thirty are collected from the statistics published in Dr. Beddoe's paper on the stature and bulk of man in the British Isles.

The heights and weights of the idiots and imbeciles are derived from the institutions enumerated at the head of the columns, and were collected for me by Drs. Grabham, Ireland, and Shuttleworth, the medical superintendents.

For the purpose of comparing together English and American born boys, I have added to the tables two columns of averages from Dr. Bowditch's report on the growth of children in Boston and the surrounding neighbourhood. As they include all classes of society and are both town and country bred, they do not correspond with any of the classes I have adopted, but it will be seen that they closely resemble the two divisions of Class I. of our English boys. I have to regret that I have not been able to include some statistics promised by my friend Dr. F. N. Manning of Sydney, to show if any changes have taken place in the physique of our Australian relations.\*

The study of the extremes of the stature which the body may attain in giants and dwarfs is full of interest and importance. Giants and dwarfs occur at all ages; but as growth is very irregular in many individuals, sometimes being excessive at an early age and occasionally retarded for years, the true giant

\* It does not form part of my present plan to point out the numerous interesting conclusions to be drawn from these tables. My object here is to show how imperfect our information is, and to endeavour to direct into proper channels the labours of many different observers. Statistics of the kind embodied in my tables, collected, apparently, without any definite object, lie scattered all over the country, and many opportunities of adding to them are lost for want of an intelligent motive. I hope that my present work will explain the object, and supply a motive for more extended efforts.

and dwarf can only be distinguished at the time of full development, unless, indeed, the proportions are very unusual. In the giant the great stature is due, chiefly, to an excessive growth of the lower extremities, the size of head and trunk being nearly the same as that of the mean boy or man of the same age. The Swedish giant was about  $8\frac{1}{4}$  feet in height, being the tallest man on record. In the natural dwarf, where the proportions are fairly uniform, the head is always large in proportion to the rest of the body, as it is in the young child; indeed, the proportions of the dwarf Tom Thumb are almost exactly those of an ordinary infant of from thirteen to fifteen months old.

TABLE XII. *Showing the average HEIGHT (without shoes) of 54,447 Boys  
4630 of America*

Male population and percentage, including children, 1871.	Professional classes.				Commercial classes.				Labouring					
	Class I.				Class II.				Class III.		Class IV.		Class V.	
	328,270 =4.46 per cent.				762,014 =10.36 per cent.				439,377 =5.97 per cent.		2,383,799 =32.41 per cent.		232,784 =3.16 per cent.	
Social position, occupation, 'nurture.'	No occupation. Officers of Army and Navy. Clergymen. Physicians and Surgeons. Barristers and Solicitors. Civil Servants. Bankers. Merchants.				Clerks. Shopkeepers.				Soldiers. Policemen. Messengers. Servants.		Labourers: Agricultural, Roads, Quarries, Navvies, Railways.		Sailors. Fishermen. Bargemen.	
Sanitary surroundings.	Outdoor. Country.		Indoor. Towns.		Indoor. Towns.		Selected.		Outdoor. Country.		Outdoor. Seafaring.			
Age last birthday. Birth.	No. obs.	Inches.	No. obs.	Inches.	No. obs.	Inches.	No. obs.	Inches.	No. obs.	Inches.	No. obs.	Inches.		
1	—	—	—	—	—	—	—	—	—	—	—	—		
2	—	—	—	—	—	—	—	—	—	—	—	—		
3	—	—	—	—	—	—	—	—	—	—	—	—		
4	—	—	—	—	—	—	—	—	—	—	—	—		
5	—	—	—	—	—	—	138	41.25	—	—	—	—		
6	—	—	—	—	—	—	287	42.97	—	—	—	—		
7	—	—	3	46.16	—	—	730	45.22	—	—	—	—		
8	—	—	16	47.31	—	—	860	46.98	133	46.91	—	—		
9	—	—	59	50.18	—	—	826	48.94	156	49.43	—	—		
10	74	53.40	160	52.73	—	—	783	50.31	143	51.23	356	51.04		
11	150	54.91	294	53.84	—	—	840	51.20	138	52.94	166	52.01		
12	248	56.97	495	55.51	—	—	426	52.52	100	55.19	20	54.15		
13	473	58.79	501	58.30	—	—	415	54.65	9	56.05	393	55.31		
14	477	61.11	508	59.84	—	—	1302	56.08	—	—	100	58.91		
15	541	63.47	303	62.60	—	—	140	59.10	—	—	—	—		
16	686	66.40	76	65.10	—	—	133	63.90	—	—	—	—		
17	1602	67.86	57	66.74	—	—	870	65.91	—	—	—	—		
18	1522	68.29	23	68.00	—	—	5246	66.78	—	—	—	—		
19	794	68.72	3	69.16	—	—	3385	67.19	—	—	—	—		
20	391	69.13	—	—	—	—	2466	67.26	—	—	—	—		
21	326	68.90	—	—	—	—	1710	67.53	—	—	—	—		
22	198	68.65	—	—	—	—	1568	67.62	—	—	—	—		
23-30	170	68.66	—	—	—	—	3256	67.93	—	—	—	—		
23-50	—	—	115	67.93	242	67.28	4144	67.22	1243	67.10	313	66.31		
—	7652	—	2613	—	242	—	29525	—	1922	—	1348	—		

*and Men of different Classes (and of Idiots) of the English Population, and of Parentage (Boston, U.S.A.).*

classes.		Industrial classes.						United States of America. Of American parentage.				Age last birthday. Birth.
Class VI.		Class VII.		Class VIII.								
435,558 = 5.92 per cent.		1,971,295 = 26.82 per cent.		801,536 = 10.9 per cent.		IDIOTS and IMBECILES: Earlswood Asylum, Royal Albert Asylum, Larbert Institution for Imbeciles.		Public and private Latin Schools, Massachusetts Institute of Technology.		Public schools of the city of Boston, South Boston, Roxbury, Charlestown, Jamaica Plain, and Brooklyn.		
Miners: Coal, Mineral.		Artisans.		Factory operatives, Sedentary trades: Tailors, Shoemakers, &c.								
Underground.		Indoor. Towns.		Indoor. Towns.				Selected.		Town and Country.		
No. obs.	Inches.	No. obs.	Inches.	No. obs.	Inches.	No. obs.	Inches. Mean height.	No. obs.	Inches.	No. obs.	Inches.	
—	—	100	19.34	—	—	—	—	—	—	—	—	1
—	—	—	—	—	—	—	—	—	—	—	—	2
—	—	—	—	—	—	—	—	—	—	—	—	3
—	—	21	38.45	—	—	3	38.00	—	—	—	—	4
—	—	37	40.10	—	—	13	40.03	—	—	201	41.74	5
—	—	40	43.27	—	—	34	42.23	—	—	342	44.10	6
—	—	53	45.70	—	—	41	44.12	—	—	369	46.21	7
—	—	110	46.88	162	46.90	54	45.74	—	—	407	48.16	8
—	—	295	48.97	276	48.46	57	47.49	2	52.00	381	50.09	9
—	—	279	50.77	419	50.21	72	49.11	19	53.51	360	52.21	10
—	—	175	52.60	3415	51.56	61	50.97	17	54.90	350	54.01	11
—	—	151	53.95	325	53.36	63	52.46	23	56.78	373	55.78	12
—	—	550	56.96	—	—	74	54.75	41	59.60	391	58.17	13
—	—	946	58.15	—	—	56	56.53	49	61.51	386	61.08	14
—	—	605	60.92	—	—	54	59.26	46	64.20	342	62.96	15
—	—	895	63.78	—	—	66	60.69	40	65.83	232	65.58	16
—	—	449	64.40	—	—	36	62.67	32	67.44	128	66.29	17
—	—	153	65.30	—	—	37	63.21	29		65	66.76	18
—	—	97	66.02	—	—	25	63.19	—	—	—	—	19
—	—	69	66.31	—	—	28	64.18	—	—	—	—	20
—	—	55	66.84	—	—	20	64.25	—	—	—	—	21
—	—	36	66.25	—	—	16	64.50	—	—	—	—	22
—	—	166	66.43	—	—	—	—	—	—	—	—	23-30
67	66.91	335	66.77	135	66.89	19	64.87	—	—	—	—	23-50
67	—	5517	—	4732	—	829	—	303	—	4327	—	



TABLE XIII. *Showing the average WEIGHT (including clothes) of 54,447 Boy  
4630 of America*

Male population and percentage, including children, 1871.	Professional classes.				Commercial classes.				Labouring					
	Class I.				Class II.				Class III.		Class IV.		Class V.	
	328,270 =4·46 per cent.				762,014 =10·36 per cent.				439, 377 =5·97 per cent.		2,383,799 =32·41 per cent.		232,784 =3·16 per cent.	
Social position, occupation, 'nurture.'	No occupation. Officers of Army and Navy. Clergymen. Physicians and Surgeons. Bartenders and Solicitors. Civil Servants. Bankers. Merchants.				Clerks. Shopkeepers.				Soldiers. Policemen. Messengers. Servants.		Labourers : Agricultural, Roads, Quarries, Navvies, Railways.		Sailors. Fishermen. Bargemen.	
Sanitary surroundings.	Outdoor. Country.		Indoor. Towns.		Indoor. Towns.		Selected.		Outdoor. Country.		Outdoor. Seafaring.			
Age last birthday. Birth.	No. obs.	Lbs.	No. obs.	Lbs.	No. obs.	Lbs.	No. obs.	Lbs.	No. obs.	Lbs.	No. obs.	Lbs.	No. obs.	Lbs.
1	—	—	—	—	—	—	—	—	—	—	—	—	—	—
2	—	—	—	—	—	—	—	—	—	—	—	—	—	—
3	—	—	—	—	—	—	—	—	—	—	—	—	—	—
4	—	—	—	—	—	—	—	—	—	—	—	—	—	—
5	—	—	—	—	—	—	133	37·31	—	—	—	—	—	—
6	—	—	—	—	—	—	287	40·67	—	—	—	—	—	—
7	—	—	3	54·83	—	—	728	45·33	—	—	—	—	—	—
8	—	—	16	60·00	—	—	860	48·78	133	54·53	—	—	—	—
9	—	—	59	62·02	—	—	824	53·56	156	61·60	—	—	—	—
10	74	67·44	160	66·34	—	—	781	56·48	143	67·03	356	69·3	—	—
11	150	72·94	294	69·66	—	—	840	59·19	138	72·78	166	74·1	—	—
12	248	80·33	495	75·12	—	—	426	63·02	100	77·28	20	77·9	—	—
13	473	88·66	501	81·94	—	—	414	70·82	9	83·83	393	88·2	—	—
14	477	99·21	508	92·24	—	—	1302	80·57	—	—	100	98·5	—	—
15	541	110·42	303	105·30	—	—	140	95·91	—	—	—	—	—	—
16	686	128·34	76	118·75	—	—	133	110·80	—	—	—	—	—	—
17	1602	141·08	57	127·50	—	—	870	124·01	—	—	—	—	—	—
18	1522	146·00	23	140·00	—	—	5246	129·43	—	—	—	—	—	—
19	794	148·20	3	141·16	—	—	3386	133·51	—	—	—	—	—	—
20	391	152·07	—	—	—	—	2466	136·49	—	—	—	—	—	—
21	326	152·34	—	—	—	—	1710	137·92	—	—	—	—	—	—
22	198	152·06	—	—	—	—	1568	139·19	—	—	—	—	—	—
23-30	170	152·00	—	—	—	—	3256	141·10	—	—	—	—	—	—
23-50	—	—	70	162·10	242	146·74	4144	138·27	1243	149·22	228	163·1	—	—
Weight of clothes (approximate).	Age 5 to 10, — 10 to 30, 9 lbs.		— 6 lbs. — 9 lbs.		— — — —		— 7 lbs. — 10 lbs.		— 7 lbs. — 10 lbs.		— 7 lbs. — 10 lbs.		— 7 lbs. — 10 lbs.	

*and Men of different Classes (and of Idiots) of the English Population, and of Parentage (Boston, U.S.A.).*

classes.		Industrial classes.				IDIOTS and IMBECILES : Earlswood Asylum, Royal Albert Asylum, Larbert Institution for Imbeciles.		United States of America. Of American parentage.				Age last birthday. Birth.
Class VI.		Class VII.		Class VIII.				Public and private Latin Schools, Massachusetts Institute of Technology.		Public schools of the city of Boston, South Boston, Roxbury, Charlestown, Jamaica Plain, and Brooklyn.		
435,558 =5.92 per cent.		1,971,295 =26.82 per cent.		801,536 =10.9 per cent.								
Miners : Coal, Mineral.		Artisans.		Factory operatives. Sedentary trades : Tailors, Shoemakers, &c.								
Underground.		Indoor. Towns.		Indoor. Towns.				Selected.		Town and Country.		
No. obs.	Lbs.	No. obs.	Lbs.	No. obs.	Lbs.	No. obs.	Lbs. Mean weight.	No. obs.	Lbs.	No. obs.	Lbs.	
—	—	100	7.55	—	—	—	—	—	—	—	—	
—	—	—	—	—	—	—	—	—	—	—	—	
—	—	—	—	—	—	—	—	—	—	—	—	
—	—	—	—	—	—	3	35.0	—	—	—	—	
—	—	—	—	—	—	13	39.0	—	—	127	39.8	
—	—	—	—	—	—	34	43.0	—	—	236	43.8	
—	—	—	—	—	—	41	46.5	—	—	346	48.0	
—	—	—	—	—	—	54	50.5	—	—	338	52.9	
—	—	110	58.78	173	54.07	57	52.5	2	60.1	323	57.5	
—	—	296	63.31	331	58.56	72	59.0	19	70.6	336	64.1	
—	—	281	68.96	411	61.55	61	64.5	17	75.3	290	70.2	
—	—	175	73.94	345	66.68	63	70.5	28	85.9	309	81.3	
—	—	146	77.64	319	70.57	74	77.0	41	94.4	307	91.9	
—	—	606	89.82	18	77.40	56	85.5	49	99.9	290	100.3	
—	—	892	96.84	—	—	54	94.5	46	116.0	255	108.4	
—	—	770	107.25	—	—	66	103.0	40	125.8	238	112.9	
—	—	895	117.51	—	—	36	110.0	32	135.2	168	115.8	
—	—	501	126.53	—	—	37	116.0	29	138.2	118	115.8	
—	—	145	133.14	—	—	25	120.5	—	—	—	—	
—	—	105	138.39	—	—	28	121.5	—	—	—	—	
—	—	—	—	—	—	20	122.0	—	—	—	—	
—	—	122	140.22	—	—	16	122.5	—	—	—	—	
—	—	39	141.13	—	—	—	—	—	—	—	—	
—	—	121	144.73	—	—	—	—	—	—	—	—	
67	148.21	335	142.52	135	144.49	19	123.0	—	—	—	—	
—	—	—	6 lbs. 9 lbs.	—	7 lbs. 10 lbs.	—	6 lbs. 9½ lbs.	At 5, 2½ lbs. At 12, 7½ lbs.	At 8, 4 lbs. At 15, 8 lbs.	—	—	

*1. Of the Proportions of the Head.*

The head is one of the most essential parts of the body, and is more completely developed at birth and varies least in its proportions during growth. Being least liable to change, it has from the earliest times been made the standard for judging of the height and other proportions of the body. In the adult it is generally considered as forming the seventh part of the whole height. The proportion may, however, vary between six and eight, and in the case of giants, nine times; while in dwarfs it may form a fourth part of the height. At other periods of life it is not available as a standard, as the head and the other portions of the body do not increase in the same proportions. From birth to the period of full development the head only doubles its height, while the whole body elongates three or four times its original dimensions.

The following table (from Quetelet) shows the height of the head, and the ratio between the height of the head and that of the whole body, at different ages, in both sexes :

Age.	Men.			Women.		
	Total height.	Height of head.	Ratio.	Total height.	Height of head.	Ratio.
Birth.	Inches. 19·68	Inches. 4·37	4·50	Inches. 19·44	Inches. 4·37	4·45
1 year.	27·48	6·06	4·53	27·16	6·06	4·48
2 years.	31·14	6·81	4·57	30·74	6·77	4·54
3 "	34·03	7·16	4·74	33·64	7·08	4·74
5 "	38·86	7·56	5·14	38·34	7·40	5·18
10 "	50·11	8·07	6·21	49·37	7·91	6·21
15 "	59·56	8·46	7·04	58·58	8·38	6·99
20 "	65·74	8·93	7·35	61·96	8·66	7·15
30 "	66·37	8·97	7·39	62·20	8·80	7·15
40 "	66·37	8·97	7·39	62·20	8·80	7·15

From these figures it will be seen that the height of the head varies little in the two sexes, and it is at the time of birth half what it will attain to after the complete development of the individual. The increase is greatest during the first and second year, when it amounts to 2·44 inches. At the age of adolescence growth has almost terminated, and at the full period of development it has attained nearly 9 inches. It would appear, from M. Quetelet's observations, that the lower parts of the face grow at a greater rate than the upper; for the nasal incision which, after the age of puberty, divides the face into two equal parts, is in infants nearer the chin than the vertex.

The antero-posterior diameter of the head is a little greater at birth than the transverse, and this ratio is preserved through life. The difference between the diameter and height of the head at birth is about

1.73 inches, but in adult life this is reduced to half an inch. All the horizontal measurements of the head develop less than those of height; the ratio of those of height being as 1 to 2, the transverse as 2 to 3.

## 2. *Of the Neck.*

The growth of the neck is nearly the same as that of the head, but is less than that of the trunk, and is more perceptible in man than in woman. At birth the height of the neck is about an inch, but on account of the plumpness of the infant's chin it shortens, and it is not till the sixth or seventh year that it disengages itself and commences to grow perceptibly. After adolescence its height is about 2 inches. The ratio of growth in man is as 1 to 2; in woman as 1 to 1.79.

The diameter of the neck taken a little above the clavicles is about 1.81 inches at birth. It develops rapidly during the first year, and at the age of six years it is about  $2\frac{1}{4}$  inches for both sexes. At the age of puberty the growth becomes more rapid, especially in women; and when development has terminated both diameters of the neck are about  $4\frac{3}{4}$  inches. The circumference of the neck immediately below the chin grows in the ratio of 1 to 2.31 in men, and of 1 to 2.09 in women.

### 3. *Of the Trunk.*

The trunk is the part of the body which is included between the clavicles above and a horizontal line passing under the perineum or fork; and consists of the *chest*, the *abdomen*, and the *pelvis*.


We have seen that the height of the head increases, progressively and at the period of full development is doubled; the height of the neck increases less regularly at first, but ultimately it attains double its original dimensions. The other parts of the body increase with greater energy, and we find that the growth is greater the further the parts are situated from the summit of the head. Thus, while the measurements of the head and neck are only doubled, those of the trunk are tripled, and those of the lower extremities are more than quadrupled.

The *diameters* of the trunk, with respect to *width*, grow nearly in the same ratio as the height; they triple from birth to the period of full development. At the age of six or seven years the diameters at the shoulders, the chest, and the hips at the trochanters are doubled. The *diameters* of the trunk with respect to *thickness* grow less rapidly. The diameter of the chest from front to back only doubles towards the age of puberty, and from birth it grows only in the ratio of 1 to 2·36.

The *circumference* of the trunk grows in almost the same proportion as the height and transverse diameters. At puberty the circumferences of the trunk are much modified by sex; the shoulders are proportionably broader in man and the hips in woman, though the difference is much less than is generally believed. The development of the breasts in woman renders the chest-girth exceedingly variable, as does also her costume the circumference of the waist.


From the great importance to life of the organs contained within the chest, the circumference of that portion of the trunk has been selected, together with the total height and weight of the body, to determine the physical capacity of the individual for military or other duties. Tables III. and VIII. give the results of the measurement of the *empty* chest of males from the age of 5 to 30 years.

At the time of birth, when the child is about the sixth of the height it will ultimately attain to, the point which divides the total height into two equal parts is a little above the navel; at two years of age it is at the navel; at three years, when the child has attained half its total height, the central point is on a line with the upper borders of the iliac bones; at ten years of age, when the child has attained three-fourths of its total height, the central point is on a line with the trochanters; at thirteen years it is at






III. Diagram showing the development of  $t$   
(the black lines) and in Circum



nd 14                      10                      20                      30                      inches 40





the pubes, and in the adult man it is nearly half an inch lower. In the adult woman the central point is a little above the pubes. Proportions being kept, the distances from the vertex to the various parts of the trunk are greater in woman than in man.

#### 4. *Of the Upper Extremities.*

The upper extremity extends from the acromial end of the clavicle (the bony prominence of the shoulder) to the tip of the middle finger, and consists of three portions—the *arm*, comprised between the acromion projection and the external condyle of the humerus (the bony prominences above the elbow); the *forearm*, extending from the points just mentioned to the styloid process of the radius at the root of the thumb; and the *hand*.

It is generally believed that the space covered by the arms extended horizontally is equal to the total height of the man; but this is true only within certain limits, namely, from the time of birth to that of puberty; and this is equally true of both sexes, the difference, indeed, being less in girls than in boys. After puberty more decided changes in the proportions take place, the horizontal being greater than the perpendicular measurement, especially in men, whose chest and shoulders have a greater development in breadth than women. The ratio of height

to the measurement of the extended arms is in the adult man as 1 to 1·045, and in woman as 1 to 1·015.

The length of the arm, without including the hand, is doubled at the age between four and five years, tripled between thirteen and fourteen years, and quadrupled at the period of full development. The *hand* develops less rapidly. Its length is doubled between five and seven years, and tripled at adult age. The length of the hand, like the height of the head and the length of the foot, has been employed by artists, sculptors, and authors as a standard of the proportions of the body, its length having the ratio to the total height of the individual of 1 to 9 from about the age of seven or eight years to adult life; before that age the hand is comparatively a little longer. These proportions are truer for the woman than the man. In proportion to the length, the hand is a little broader in women.

The *forearm* is the part which increases most sensibly in length; in the new-born infant it measures about  $2\frac{1}{4}$  inches, and in the adult  $9\frac{1}{2}$  inches, the growth being in the ratio of 1 to 4·26. The 'cubit,' or the length of the forearm of the adult, was the unit of measurement among the ancients.

The *arm* (between the shoulder and elbow) measures at birth  $3\frac{1}{2}$  inches, and in the adult 13 inches; these numbers are in the ratio of 1 to 3·78.

The circumferences at the biceps and at the elbow increase nearly at the same rate, but more rapidly than the hand, viz. from 1 to  $2\frac{3}{4}$  nearly. The circumference of the biceps and the thickest part of the forearm is about the same in the adult; but both these measurements are liable to great variation under the influence of disease, exercise, &c.

#### 5. *Of the Lower Extremity.*

The lower extremity includes the *thigh*, the *leg*, and the *foot*, and is joined to the trunk on a line with the trochanters. To determine the share it takes in forming the *total height* of the individual, it is necessary to measure the distance from the perineum, or fork, to the sole of the foot; but as the relative position of the perineum is liable to vary, from obesity and other causes, measurements for determining the size of the *limb* should be made from the trochanters to the sole. The division between the thigh and the leg is at the lower edge of the patella or knee-cap, and between the leg and the foot at the bony prominences above the ankle joint—the *malleoli*.

It has been already observed that the lower extremities develop more rapidly and in greater relative proportions than other parts of the body. Thus from birth to maturity, while the head and neck double their height, and the trunk increases to three times

its original proportions, the lower extremities in the adult are five times the length they were in the newborn infant. The upper extremities grow less rapidly, the arms of the adult man being only four times the length they were at birth. Measuring from the fork to the sole of the foot, the lower limbs double their length before the third year; at twelve years they are four times, and at twenty years five times, their original length. The thigh, the leg, and the foot do not increase in the same proportions.

The length of the *thigh*, measuring from the fork to the patella or knee-cap, is  $1\frac{3}{4}$  inches in the newborn child, and nearly 13 inches in the fully developed man. These numbers are in the ratio of 1 to 7·31, and this portion of the limb acquires seven times its primitive length, an increase much greater than takes place in any other part of the body. The length of the thigh, measured from the trochanter to the knee-cap, is at birth 3·15 inches, and in the adult 15·75 inches—numbers having the ratio of 1 to 5. The length of this part of the body varies greatly in different individuals, and on it depends much of the inequality of their total heights.

The length of the *leg*, measuring from the lower edge of the patella to the inner malleolus at the ankle, is 3·42 inches at birth, and 15·35 inches in the adult, an increase in the ratio of 1 to  $4\frac{1}{2}$ .

The *height* of the foot, which is about an inch at birth, is  $3\frac{1}{4}$  inches in man fully developed, an increase in the ratio of about 1 to 3. The *length* of the foot, from the heel to the extremity of the great toe, grows a little more rapidly than the height, the ratio of increase being as 1 to  $3\frac{1}{2}$ .

From these measurements it results that the thigh grows more than the leg, and the leg than the foot. Something similar was observed in the upper extremity, the forearm, however, having the greater growth. The *circumference* of the leg at the calf is nearly the same as that at the knee.

The *foot*, like the hand and the head, has been employed as a standard of the proportions of the body. At all ages of life and in both sexes it forms from the 0.15 to 0.16 of the total height of the individual; it is, however, comparatively a little longer at the period of adolescence, but rather shorter in children and adults. Taking the length of the foot for unity, the total height of man would be  $6\frac{3}{4}$ , and of women  $6\frac{1}{4}$ . It is generally believed that the length of the foot is equal to the height of the head; but this is only true of the age of ten years: before that period the head is longer, and after it shorter, than the foot. It is also said that the length of the foot is equal to the circumference of the fist; but



there appears to be no more foundation for this than the former opinion.

NOTE. The standards of the proportions of the body employed by ancient and Renaissance sculptors and artists were taken from different parts of the body; and although they are not of much value to science, they are full of interest to those who appreciate their incomparable works of art. They were :

The *cubit* of the Egyptians, or the distance between the elbow and the extremity of the fingers; it forms the fourth part of the height of man.

The *foot*, which forms the sixth part.

The *head*, which, according to Vitruvius, forms the eighth part; but, properly speaking, the head is contained seven and a half times in the height.

The *face* (*volto*), which is equal to the length of the hand, and is the ninth part of the total height.

# ANTHROPOMETRY.

*A List of some Anthropometrical Apparatus made by*  
HAWKSLEY, 300 Oxford Street, London, W.

**STANDARD WEIGHING APPARATUS**, weighing from 2 oz. to 20 st. £5 5s.

The *Lancet*, December 7, 1872, says: 'Mr. H. has brought out a weighing-machine which is both satisfactory in its working and comparatively low in price.' At the request of Mr. F. GALTON, that the Anthropological Society might be furnished with the age, height, and weight of every boy at Marlborough College, this office was undertaken by Dr. Fergus, who says: 'The weight was taken by one of Hawksley's weighing-machines, which, for simplicity and accuracy of action, cannot be surpassed. After all the weighing was done, it had not varied the fraction of an ounce' (vide *Marlborough College Report*, 1874). This apparatus is easily taken to pieces for the convenience of packing, requires no oil, will bear severest usage, and has two adjustments, which enable any one to correct it. The profession are informed that an accurate register is kept at the above address of the weights and heights of patients, who are also supplied with a Reference Card at a charge of 1s. (Vide page 33 of this Work for illustrations.)

The above apparatus, fitted with a stadiometer (or height measurer), divided into feet, inches, and tenths, or centimètres and millimètres, or both, upon one staff, £7 7s.

**SIMPLE STADIOMETER**, consisting of a cubical rod of boxwood 2 mètres in length, with sliding rectangular bar, fitted with spirit-level, and divided into feet, inches, and tenths of an inch; also with centimètres and millimètres. 35s.

Ditto, with only 1 scale, 30s.; ditto, without spirit-level, 25s.; ditto, with 2 scales, spirit-level, and on heavy foot, 50s.

**CHAIR**, with back and seat forming a rectangle, fitted with stadiometer to measure the height of the patient standing erect, and also his sitting height. £4 4s.

**CYRTOMETER** for accurately obtaining the circumference of the chest, and also its contour. In simple form. By Hawksley. 7s. 6d.

Ditto, with improved hinge to fit the spine, and clamp in front. After Dr. Greenfield. 12s. 6d.

**DOUBLE CHEST TAPE**, reading from the spine to the right and left. By Hawksley. 5s. Strips of flexible metal for obtaining the contour of pelvis, thorax, head, &c.; or the outlines of limbs, &c.

**COXANKYLOMETER**, a T-shaped instrument for measuring the relative length of the legs in hip-disease, or from other causes. The vertical leg of the T is placed between the legs of the patient when lying upon a hard couch or table; the cross at the top is then brought level, or even, with the anterior superior spines of the ileum. If there be any difference in the length of the limbs, a comparison may be at once made and measured. In simple form. 7s. 6d.

Ditto, of more elaborate form and greater usefulness and accuracy, 21s.

**INSTRUMENT FOR MEASURING** the Angles of Limbs and their Circumferences. After Dr. Russell Reynolds. The circumference of a limb varies from position, contraction, &c., of a muscle, and the degree of tension by which it is encircled by the measuring-tape. With this instrument the angle at which a limb is measured may be recorded by an arc; the point at which it is measured is determined by a scale of inches; the circumference is measured by a tape attached to a scale of parts or degrees of tension: so that, after noting the above four indications, the limb may be examined at a future time under very precisely similar conditions. Useful in medico-legal investigations. £3 3s.

**DYNAMOGRAPH**. This is an adaptation of Duchenne's instrument to a recording plate. The variations in sustained muscular contraction, in a given space of time, are portrayed by a wave or zigzag line. £7 7s.

**DYNAMOMETER** for showing the strength of the back, or back and legs combined, in lifting from the ground.

**DYNAMOMETER** for indicating in pounds the force of a blow of the fist.

**DYNAMOMETER** of Duchenne. 35s. This instrument indicates and registers the force exerted by the grasp of the hand in kilogrammes, or in pounds avoirdupois. It is also used for exhibiting the pulling or tearing power of the two arms, the hands being attached to the opposite ends of the instrument.

**DYNAMOMETER** of Dr. Gowers. This instrument consists of a hollow steel cylinder divided on one side, about 2 inches in diameter and 4 inches in length

A dial with pointer and register is fixed at one end to indicate the degree of compression exerted by the patient. £3 3s.

NOTE. There has not yet been invented a dynamometer which gives even an approximate of the power of compression exerted by the hand. As in spirometers of lung capacity a 'knowing one' will be able to give unusual results, so a particular and often a accidental grasp of any present hand dynamometer will give unreliable results.

**DYNAMOMETER** for testing the combined strength of the muscles of the arms and back. The position of testing with this instrument is exactly like that of drawing a bow in archery.

**ÆSTHESIOMETER** for ascertaining the sensitiveness of the cutaneous nerves. After Dr. Ogle. With dial indicating to  $\frac{1}{100}$ th of an inch.

Ditto, with sliding points, divided into millimètres, 7s. 6d., 10s. 6d. Compasses for above purpose, 3s. 6d., 1s.

**CHEST CALLIPERS**, in boxwood, with arc divided into inches and into centimètres. 7s. 6d.

Ditto, ditto, in steel; very accurate; with arc and clamping screw capable of measuring to 15 inches diameter, 35s.

**STETHOGRAPH**. After Dr. Sanderson. Consisting of a rectangular frame, to one arm of which is fitted a tambour, and to the other a sliding rod to rest upon the spine. This instrument records the movement of respiration and cardiac impulse upon the recording cylinder. 63s.

**STETHOMETER** of Dr. Ransome, recording the movements upwards, outwards, and forwards of the chest in  $\frac{1}{100}$ th of an inch. £7 7s.

**STETHOMETER** of Dr. Ransome. Applied by Hawksley to measuring the upward, outward, and forward movement of every rib for the whole circumference of the thorax. With spinal rod and adjustable rods to keep the spine absolutely in one fixed position. Only one instrument has been made, and that for the late Dr. Sargent.

**RANSOME'S COMPOUND LEVER STETHOGRAPH** for recording the forward and upward movement of the thorax. The lever writes upon marked paper or glass, or with a pencil, the above movements. £5 5s.

**RANSOME'S 2-DIAL STETHOMETER** for upward and forward movements, registering to  $\frac{1}{100}$ th of an inch. On stand. £3 3s.

**INSTRUMENT** for measuring the degree of pressure required to control a hernia. With adjustable pressure screw and scale for showing in pounds and ounces the pressure exerted. The instrument half encircles the body, and has a pad at each end. One rests upon the spine, and the other is attached to a strong spring with dividing scale. A rack and pinion increases or diminishes the pressure upon the hernial opening. The degree of impulse on coughing may be measured, and also the degree of pressure required to control the protrusion of the hernia. After Hawksley. A compact name is lacking for this instrument.

**SPIROMETER**. After Lowne. This instrument is very portable and sensitive. £5 10s.

Ditto. After Hutchinson. This instrument accurately determines the capacity of the chest. £5.

**PERIMETERS** and **CHARTS** for measuring and mapping the sensitive field of the retina. £3 3s.

**IVORY SCALES** for the Pocket. Containing the centigrade and Fahrenheit temperature scales side by side for instant comparison; also centimètres divided into millimètres, and English inches divided into  $\frac{1}{10}$ ths,  $\frac{1}{20}$ ths, and  $\frac{1}{40}$ ths. 1s. 6d.

**OCCIPITAL CROTCHET** of M. Broca.

**STEREOGRAPH** and **CRANIOPHERE** of M. Broca.

**CRANIOPHERE**. After Topinhard.

**PARIETAL GONIOMETER**. By Quatrefages.

**SPHENOIDAL CROTCHET**. After M. Broca.

**SCALE** (Sliding and Rectangular) for taking measurements of the projections of the head in the living subject.

**GONIOMETER** for measuring the median facial angles. M. Broca.

**INSTRUMENT** for testing the higher limits of audition. After F. Galton.

**INSTRUMENT** for measuring the rapidity of perception, consisting of a graduated board, at each end of which are clamps and arrangements for releasing and stopping a cord which is pulled by a falling weight. The signal indicating to the patient that the weight has begun to fall is by a bell or sharp tap, or by both. Upon hearing this sound, the patient immediately compresses one of the clamps beneath his fingers, and stops the falling weight; then the distance which the weight has fallen is indicated by the cord on the divided board in  $\frac{1}{100}$ th or  $\frac{1}{1000}$ th of a second. After F. Galton.



*London, New Burlington Street.  
July, 1878.*

SELECTION  
FROM  
MESSRS J. & A. CHURCHILL'S  
*General Catalogue*  
COMPRISING  
ALL RECENT WORKS PUBLISHED BY THEM  
ON THE  
ART AND SCIENCE  
OF  
M E D I C I N E

# INDEX

	PAGE		PAGE
Acton on the Reproductive Organs . . .	8	Cullingworth's Nurse's Companion . . .	14
Adams (W.) on Clubfoot . . .	6	Curling's Diseases of the Rectum . . .	7
— (R.) on Rheumatic Gout . . .	18	Dalby on the Ear . . .	6
Allen on Aural Catarrh . . .	6	Dalton's Human Physiology . . .	9
Allingham on Diseases of Rectum . . .	8	Day on Children's Diseases . . .	13
Anatomical Remembrancer . . .	11	— on Headaches . . .	18
Anderson (McC.) on Eczema . . .	19	De Valcourt on Cannes . . .	16
— (McC.) on Parasitic Affec- tions . . .	19	Dobell's Lectures on Winter Cough . . .	15
Arnott on Cancer . . .	19	— First Stage of Consumption . . .	15
Aveling's English Midwives . . .	14	Domville's Manual for Hospital Nurses . . .	14
Balfour's Diseases of the Heart . . .	16	Druitt's Surgeon's Vade-Mecum . . .	4
Bantock's Rupture of Perineum . . .	14	Dunglison's Medical Dictionary . . .	22
Barclay's Medical Diagnosis . . .	11	Ellis's Manual of Diseases of Children . . .	12
Barker's Puerperal Diseases . . .	13	Fayrer's Observations in India . . .	4
Barnes' Obstetric Operations . . .	14	Fergusson's Practical Surgery . . .	4
— Diseases of Women . . .	14	Fenwick's Guide to Medical Diagnosis . . .	11
Basham on Renal Diseases . . .	8	— on the Stomach, &c. . .	18
— on Diseases of the Kidneys . . .	8	Flint on Phthisis . . .	15
Beale on Kidney Diseases . . .	8	— on Percussion and Auscultation . . .	15
— on Microscope in Medicine . . .	11	Flower's Nerves of the Human Body . . .	10
Bellamy's Guide to Surgical Ana- tomy . . .	10	Foster's Clinical Medicine . . .	12
Bennet's Winter and Spring on the Mediterranean . . .	16	Fox (T.) Atlas of Skin Diseases . . .	19
— Pulmonary Consumption . . .	16	Fox and Farquhar's Skin Diseases of India . . .	20
— Nutrition . . .	18	Frey's Histology . . .	9
Bennett (J. R.) on Cancerous Growths . . .	19	Gamgee on Fractures of the Limbs . . .	4
Berkart's Asthma . . .	15	— on Treatment of Wounds . . .	4
Bigg's Orthopraxy . . .	6	Gant's Science and Practice of Surgery . . .	4
Binz's Elements of Therapeutics . . .	12	— Diseases of the Bladder . . .	8
Black on the Urinary Organs . . .	8	Gaskoin on Psoriasis or Lepra . . .	19
Bose's Rational Therapeutics . . .	11	Glenn's Laws affecting Medical Men . . .	20
— Recognisant Medicine . . .	11	Godlee's Atlas of Human Anatomy . . .	11
Braune's Topographical Anatomy . . .	11	Gordon on Fractures . . .	6
Brodhurst's Orthopædic Surgery . . .	6	Habershon on Diseases of the Liver . . .	17
Bryant's Practice of Surgery . . .	4	— on Diseases of the Stomach . . .	17
Bucknill and Tuke's Psychological Medicine . . .	21	— on the Pneumogastric Nerve . . .	17
Burdett's Cottage Hospital . . .	15	Hancock's Surgery of Foot and Ankle . . .	6
Burnett on the Ear . . .	6	Harris on Lithotomy . . .	7
Buzzard on Syphilitic Nervous Affec- tions . . .	8	Harrison's Stricture of Urethra . . .	7
Carpenter's Human Physiology . . .	10	Hayden on the Heart . . .	16
Carter on Mycetoma . . .	20	Heath's Minor Surgery and Bandaging . . .	5
Cauty on Diseases of the Skin . . .	20	— Diseases and Injuries of Jaws . . .	5
Chapman on Neuralgia . . .	18	— Operative Surgery . . .	5
Charteris' Practice of Medicine . . .	11	— Practical Anatomy . . .	10
Clark's Outlines of Surgery . . .	4	Higgins' Ophthalmic Practice . . .	22
— Surgical Diagnosis . . .	5	Holden's Landmarks . . .	10
Clay's Obstetric Surgery . . .	13	Holt on Stricture of the Urethra . . .	7
Cobbold on Worms . . .	20	Hood on Gout, Rheumatism, &c. . .	18
Coles' Dental Mechanics . . .	23	Hooper's Physician's Vade-Mecum . . .	11
Cooper's Surgical Dictionary . . .	4	Horton's Tropical Diseases . . .	17
Cormack's Clinical Studies . . .	12	Hutchinson's Clinical Surgery . . .	5
Cottle's Hair in Health and Disease . . .	28	Huth's Marriage of Near Kin . . .	9
Coulson on Syphilis . . .	9	Ireland's Idiocy and Imbecility . . .	21
— on Stone in the Bladder . . .	9	James' Sore Throat . . .	16
		Jones (C. H.) and Sieveking's Patho- logical Anatomy . . .	10
		— (C. H.) on Functional Nervous Disorders . . .	18

	PAGE		PAGE
Jones (H. McN.) Aural Surgery . . . . .	6	Smith (J.) Dental Anatomy . . . . .	23
Jones (Wharton) Ophthalmic Medi- cine and Surgery . . . . .	23	Smith (W. R.) Nursing . . . . .	14
Jordan's Surgical Inflammations . . . . .	6	Spender's Bath Waters . . . . .	16
— Surgical Inquiries . . . . .	6	Steiner's Diseases of Children . . . . .	13
Leber and Rottenstein's Dental Caries . . . . .	23	Stowe's Toxicological Chart . . . . .	20
Lee (H.) Practical Pathology . . . . .	8	Sullivan's Tropical Diseases . . . . .	17
— on Syphilis . . . . .	8	Swain's Surgical Emergencies . . . . .	5
Leared on Imperfect Digestion . . . . .	18	Swayne's Obstetric Aphorisms . . . . .	14
Liebreich's Atlas of Ophthalmoscopy . . . . .	22	Taft's Operative Dentistry . . . . .	23
Living on Megrim, &c. . . . .	18	Tait's Hospital Mortality . . . . .	14
Macdonald's Examination of Water . . . . .	21	Taylor's Principles of Medical Juris- prudence . . . . .	20
Mackenzie on Growths in the Larynx . . . . .	15	— Manual of Medical Juris- prudence . . . . .	20
Macnamara on Diseases of the Eye . . . . .	22	— Poisons in relation to Medical Jurisprudence . . . . .	20
Madden's Health Resorts . . . . .	17	Thompson's Stricture of Urethra . . . . .	7
Marsden on certain Forms of Cancer . . . . .	19	— Practical Lithotomy and Lithotripsy . . . . .	7
Mason on Harelip and Cleft Palate . . . . .	5	— Diseases of Urinary Organs . . . . .	7
Mauder's Operative Surgery . . . . .	4	— Diseases of the Prostate . . . . .	7
— Surgery of Arteries . . . . .	4	— Calculous Disease . . . . .	7
Mayne's Medical Vocabulary . . . . .	22	Thornton on Tracheotomy . . . . .	16
Meryon's System of Nerves . . . . .	18	Thorowgood on Asthma . . . . .	15
Moore's Family Medicine for India . . . . .	17	— on Materia Medica . . . . .	12
Ogston's Medical Jurisprudence . . . . .	20	Thudichum's Pathology of Urine . . . . .	9
Parkes' Manual of Practical Hygiene . . . . .	21	Tibbitts' Medical Electricity . . . . .	22
Parkin's Epidemiology . . . . .	23	— Map of Motor Points . . . . .	22
Pavy on Food and Dietetics . . . . .	18	Tilt's Uterine Therapeutics . . . . .	13
Peacock's Valvular Disease . . . . .	15	— Change of Life . . . . .	13
Phillips' Materia Medica . . . . .	12	— Health in India . . . . .	17
Pirrie's Surgery . . . . .	4	Tomes' (C. S.) Dental Anatomy . . . . .	23
Pollock's Rheumatism . . . . .	19	— (J. and C. S.) Dental Surgery . . . . .	23
Ramsbotham's Obstetrics . . . . .	13	Tufnell's Internal Aneurism . . . . .	7
Reynolds' Uses of Electricity . . . . .	22	Tuke on the Influence of the Mind upon the Body . . . . .	21
Roberts' Practice of Midwifery . . . . .	13	Van Buren on Diseases of the Genito- Urinary Organs . . . . .	9
Roussel's Transfusion of Blood . . . . .	5	Veitch's Handbook for Nurses . . . . .	14
Routh's Infant Feeding . . . . .	12	Virchow's Post-mortem Examinations . . . . .	10
Roy's Burdwan Fever . . . . .	17	Wagstaffe's Human Osteology . . . . .	10
Royle and Harley's Materia Medica . . . . .	12	Walton's Diseases of the Eye . . . . .	22
Rutherford's Practical Histology . . . . .	9	Ward on Affections of the Liver . . . . .	17
Sabben and Browne's Handbook of Law and Lunacy . . . . .	21	Waring's Practical Therapeutics . . . . .	12
Salts' Medico-Electric Apparatus . . . . .	22	— Bazaar Medicines of India . . . . .	17
Sanderson's Physiological Handbook . . . . .	9	Waters on Diseases of the Chest . . . . .	15
Sansom's Diseases of the Heart . . . . .	16	Wells (Soelberg) on Diseases of the Eye . . . . .	23
Savage on the Female Pelvic Organs . . . . .	5	— Long, Short, and Weak Sight . . . . .	23
Savory's Domestic Medicine . . . . .	14	Wells (Spencer) on Diseases of the Ovaries . . . . .	13
Sayre's Orthopædic Surgery . . . . .	7	Wilks' Diseases of Nervous System . . . . .	18
Schroeder's Manual of Midwifery . . . . .	13	— Pathological Anatomy . . . . .	10
Semple on the Heart . . . . .	15	Wilson's (E.) Anatomist's Vade- Mecum . . . . .	11
Sewill's Dental Anatomy . . . . .	23	— Diseases of the Skin . . . . .	19
Shapter's Diseases of the Heart . . . . .	16	— Lectures on Ekzema . . . . .	19
Shaw's Medical Remembrancer . . . . .	12	— Lectures on Dermatology . . . . .	19
Sheppard on Madness . . . . .	21	Wilson's (G.) Handbook of Hygiene . . . . .	21
Sibson's Medical Anatomy . . . . .	10	Woodman & Tidy's Forensic Medicine . . . . .	21
Sieveking's Life Assurance . . . . .	21		
Smith (E.) Wasting Diseases of Children . . . . .	12		
— Clinical Studies . . . . .	12		
Smith (Henry) Surgery of the Rectum . . . . .	8		
Smith (Heywood) Gynæcology . . . . .	13		

**THE PRACTICE OF SURGERY :**

a Manual by THOMAS BRYANT, F.R.C.S., Surgeon to Guy's Hospital.  
Second Edition, 2 vols., crown 8vo, with 559 Engravings, 25s. [1876]

**THE PRINCIPLES AND PRACTICE OF SURGERY,**

by WILLIAM PIRRIE, F.R.S.E., Professor of Surgery in the University  
of Aberdeen. Third Edition, 8vo, with 490 Engravings, 28s. [1873]

**A SYSTEM OF PRACTICAL SURGERY,**

by Sir WILLIAM FERGUSSON, Bart., F.R.C.S., F.R.S. Fifth Edition,  
8vo, with 463 Engravings, 21s. [1870]

**OPERATIVE SURGERY,**

by C. F. MAUNDER, F.R.C.S., Surgeon to the London Hospital.  
Second Edition, post 8vo, with 164 Engravings, 6s. [1873]

BY THE SAME AUTHOR.

**SURGERY OF THE ARTERIES :**

Lettsomian Lectures for 1875, on Aneurisma, Wounds, Hæmorrhages,  
&c. Post 8vo, with 18 Engravings, 5s. [1876]

**THE SURGEON'S VADE-MECUM,**

a Manual of Modern Surgery, by ROBERT DRUITT. Eleventh Edition,  
fcap. 8vo, with 369 Engravings, 14s. [1873]

**THE SCIENCE AND PRACTICE OF SURGERY :**

a complete System and Textbook by F. J. GANT, F.R.C.S., Senior Sur-  
geon to the Royal Free Hospital. 8vo, with 470 Engravings, 24s. [1871]

**OUTLINES OF SURGERY AND SURGICAL PATHOLOGY,**

including the Diagnosis and Treatment of Obscure and Urgent  
Cases, and the Surgical Anatomy of some Important Structures and  
Regions, by F. LE GROS CLARK, F.R.S., Consulting Surgeon to St.  
Thomas's Hospital. Second Edition, Revised and Expanded by the  
Author, assisted by W. W. WAGSTAFFE, F.R.C.S., Assistant-Surgeon  
to St. Thomas's Hospital. 8vo, 10s. 6d. [1873]

**CLINICAL AND PATHOLOGICAL OBSERVATIONS IN INDIA,**

by Sir J. FAYRER, K.C.S.I., M.D., F.R.C.P. Lond., F.R.S.E., Honorary  
Physician to the Queen. 8vo, with Engravings, 20s. [1873]

**DICTIONARY OF PRACTICAL SURGERY**

and Encyclopædia of Surgical Science, by SAMUEL COOPER. New  
Edition, by SAMUEL A. LANE, Consulting Surgeon to St. Mary's and  
to the Lock Hospitals; assisted by various Eminent Surgeons. 2 vols.  
8vo, 50s. [1861 and 1873]

**TREATMENT OF WOUNDS :**

Clinical Lectures, by SAMPSON GAMGEE, F.R.S.E., Surgeon to the  
Queen's Hospital, Birmingham. Crown 8vo, with Engravings, 5s. [1873]

BY THE SAME AUTHOR,

**FRACTURES OF THE LIMBS**

and their Treatment. 8vo, with Plates, 10s. 6d. [1871]

**SURGICAL EMERGENCIES**

together with the Emergencies attendant on Parturition and the Treatment of Poisoning: a Manual for the use of General Practitioners, by WILLIAM P. SWAIN, F.R.C.S., Surgeon to the Royal Albert Hospital, Devonport. Second Edition, post 8vo, with 104 Engravings, 6s. 6d. [1876]

**TRANSFUSION OF HUMAN BLOOD:**

with Table of 50 cases, by Dr. ROUSSEL, of Geneva. Translated by CLAUDE GUINNESS, B.A. With a Preface by SIR JAMES PAGET, Bart. Crown 8vo, 2s. 6d. [1877]

**ILLUSTRATIONS OF CLINICAL SURGERY,**

consisting of Coloured Plates, Photographs, Woodcuts, Diagrams, &c., illustrating Surgical Diseases, Symptoms and Accidents; also Operations and other methods of Treatment. By JONATHAN HUTCHINSON, F.R.C.S., Senior Surgeon to the London Hospital. In Quarterly Fasciculi. Fasc. I to XI already issued. 6s. 6d. each. Fasciculi I to X bound, with Appendix and Index, £3 10s. [1876-8]

**PRINCIPLES OF SURGICAL DIAGNOSIS**

especially in Relation to Shock and Visceral Lesions, by F. LE GROS CLARK, F.R.C.S., Consulting Surgeon to St. Thomas's Hospital. 8vo, 10s. 6d. [1870]

**MINOR SURGERY AND BANDAGING:**

a Manual for the Use of House-Surgeons, Dressers, and Junior Practitioners, by CHRISTOPHER HEATH, F.R.C.S., Surgeon to University College Hospital, and Holme Professor of Surgery in University College. Fifth Edition, fcap 8vo, with 86 Engravings, 5s. 6d. [1875]

BY THE SAME AUTHOR,

**INJURIES AND DISEASES OF THE JAWS:**

JACKSONIAN PRIZE ESSAY. Second Edition, 8vo, with 164 Engravings, 12s. [1872]

BY THE SAME AUTHOR.

**A COURSE OF OPERATIVE SURGERY:**

with 20 Plates drawn from Nature by M. LÉVEILLÉ, and coloured by hand under his direction. Large 8vo. 40s. [1877]

**HARE-LIP AND CLEFT PALATE,**

by FRANCIS MASON, F.R.C.S., Surgeon and Lecturer on Anatomy at St. Thomas's Hospital. With 66 Engravings, 8vo, 6s. [1877]

**THE FEMALE PELVIC ORGANS,**

their Surgery, Surgical Pathology, and Surgical Anatomy, in a Series of Coloured Plates taken from Nature: with Commentaries, Notes, and Cases, by HENRY SAVAGE, M.D. Lond., F.R.C.S., Consulting Officer of the Samaritan Free Hospital. Third Edition, 4to, £1 15s. [1876]



**FRACTURES OF THE LOWER END OF THE RADIUS,**

Fractures of the Clavicle, and on the Reduction of the Recent Inward Dislocations of the Shoulder Joint. By **ALEXANDER GORDON, M.D.**, Professor of Surgery in Queen's College, Belfast. With Engravings, 8vo, 5s. [1875]

**DISEASES AND INJURIES OF THE EAR,**

by **W. B. DALBY, F.R.C.S., M.B.**, Aural Surgeon and Lecturer on Aural Surgery at St. George's Hospital. Crown 8vo, with 21 Engravings, 6s. 6d. [1875]

**AURAL SURGERY ;**

A Practical Treatise, by **H. MACNAUGHTON JONES, M.D.**, Surgeon to the Cork Ophthalmic and Aural Hospital. With 46 Engravings, crown 8vo, 5s. [1878.]

**THE EAR :**

its Anatomy, Physiology, and Diseases. A Practical Treatise, by **CHARLES H. BURNETT, A.M., M.D.**, Aural Surgeon to the Presbyterian Hospital, and Surgeon in Charge of the Infirmary for Diseases of the Ear, Philadelphia. With 87 Engravings, 8vo, 18s. [1877]

**AURAL CATARRH ;**

or, the Commonest Forms of Deafness, and their Cure, by **PETER ALLEN, M.D., F.R.C.S.E.**, late Aural Surgeon to St. Mary's Hospital. Second Edition, crown 8vo, with Engravings, 8s. 6d. [1874]

**CLUBFOOT :**

its Causes, Pathology, and Treatment; being the Jacksonian Prize Essay by **WM. ADAMS, F.R.C.S.**, Surgeon to the Great Northern Hospital. Second Edition, 8vo, with 106 Engravings and 6 Lithographic Plates, 15s. [1873]

**ORTHOPÆDIC SURGERY :**

Lectures delivered at St. George's Hospital, by **BERNARD E. BRODHURST, F.R.C.S.**, Surgeon to the Royal Orthopædic Hospital. Second Edition, 8vo, with Engravings, 12s. 6d. [1876]

**OPERATIVE SURGERY OF THE FOOT AND ANKLE,**

by **HENRY HANCOCK, F.R.C.S.**, Consulting Surgeon to Charing Cross Hospital. 8vo, with Engravings, 15s. [1873]

**THE TREATMENT OF SURGICAL INFLAMMATIONS**

by a New Method, which greatly shortens their Duration, by **FURNEAUX JORDAN, F.R.C.S.**, Professor of Surgery in Queen's College, Birmingham. 8vo, with Plates, 7s. 6d. [1870]

BY THE SAME AUTHOR,

**SURGICAL INQUIRIES.**

With numerous Lithographic Plates. 8vo, 5s. [1873]

**ORTHOPRAXY :**

the Mechanical Treatment of Deformities, Debilities, and Deficiencies of the Human Frame, by **H. HEATHER BIGG**, Associate of the Institute of Civil Engineers. Third Edition, with 319 Engravings, 8vo, 15s. [1877]

**ORTHOPÆDIC SURGERY :**

and Diseases of the Joints. Lectures by LEWIS A. SAYRE, M.D.,  
Professor of Orthopædic Surgery, Fractures and Dislocations, and  
Clinical Surgery, in Bellevue Hospital Medical College, New York.  
With 274 Wood Engravings, 8vo, 20s. [1876]

**INTERNAL ANEURISM :**

Its Successful Treatment by Consolidation of the Contents of the Sac.  
By T. JOLIFFE TUFNELL, F.R.C.S.I., President of the Royal College  
of Surgeons in Ireland. With Coloured Plates. Second Edition,  
royal 8vo, 5s. [1875]

**DISEASES OF THE RECTUM,**

by THOMAS B. CURLING, F.R.S., Consulting Surgeon to the London  
Hospital. Fourth Edition, Revised, 8vo, 7s. 6d. [1876]

**STRICTURE OF THE URETHRA**

and Urinary Fistulæ; their Pathology and Treatment: Jacksonian  
Prize Essay by Sir HENRY THOMPSON, F.R.C.S., Emeritus Professor  
of Surgery to University College. Third Edition, 8vo, with Plates,  
10s. [1869]

BY THE SAME AUTHOR,

**PRACTICAL LITHOTOMY AND LITHOTRITY ;**

or, An Inquiry into the best Modes of removing Stone from the  
Bladder. Second Edition, 8vo, with numerous Engravings. 10s. [1871]

ALSO,

**DISEASES OF THE URINARY ORGANS :**

(Clinical Lectures). Fourth Edition, 8vo, with 2 Plates and 59  
Engravings, 12s. [1876]

ALSO,

**DISEASES OF THE PROSTATE :**

their Pathology and Treatment. Fourth Edition, 8vo, with numerous  
Plates, 10s. [1873]

ALSO,

**THE PREVENTIVE TREATMENT OF CALCULOUS DISEASE**

and the Use of Solvent Remedies. Second Edition, fcap. 8vo, 2s. 6d.  
[1876]

**STRICTURE OF THE URETHRA**

and its Immediate Treatment, by BARNARD HOLT, F.R.C.S.,  
Consulting Surgeon to the Westminster Hospital. Third Edition,  
8vo, 6s. [1868]

**STRICTURE OF THE URETHRA,**

and other Diseases of the Urinary Organs, by REGINALD HARRISON,  
F.R.C.S., Surgeon to the Liverpool Royal Infirmary. With 10 plates,  
8vo, 7s. 6d. [1878.]

**LITHOTOMY AND EXTRACTION OF STONE**

from the Bladder, Urethra, and Prostate of the Male, and from the  
Bladder of the Female, by W. POULETT HARRIS, M.D., Surgeon-Major  
H.M. Bengal Medical Service. With Engravings, 8vo, 10s. 6d. [1876]

**THE SURGERY OF THE RECTUM :**

Lettsomian Lectures by **HENRY SMITH, F.R.C.S.**, Professor of Surgery in King's College, Surgeon to King's College Hospital. Fourth Edition, fcap. 8vo, 5s. [1876]

**FISTULA, HÆMORRHOIDS, PAINFUL ULCER,**

Stricture, Prolapsus, and other Diseases of the Rectum: their Diagnosis and Treatment, by **WM. A. LLINGHAM, F.R.C.S.**, Surgeon to St. Mark's Hospital for Fistula, &c. Second Edition, 8vo, 7s. [1873]

**KIDNEY DISEASES, URINARY DEPOSITS,**

and Calculous Disorders by **LIONEL S. BEALE, M.B., F.R.S., F.R.C.P.**, Physician to King's College Hospital. Third Edition, 8vo, with 70 Plates, 25s. [1868]

**DISEASES OF THE BLADDER,**

Prostate Gland and Urethra, including a practical view of Urinary Diseases, Deposits and Calculi, by **F. J. GANT, F.R.C.S.**, Senior Surgeon to the Royal Free Hospital. Fourth Edition, crown 8vo, with Engravings, 10s. 6d. [1876]

**RENAL DISEASES :**

a Clinical Guide to their Diagnosis and Treatment by **W. R. BASHAM, M.D., F.R.C.P.**, late Senior Physician to the Westminster Hospital. Post 8vo, 7s. [1870]

BY THE SAME AUTHOR,

**THE DIAGNOSIS OF DISEASES OF THE KIDNEYS,**

with Aids thereto. 8vo, with 10 Plates, 5s. [1873]

**THE REPRODUCTIVE ORGANS**

in Childhood, Youth, Adult Age, and Advanced Life (Functions and Disorders of), considered in their Physiological, Social, and Moral Relations, by **WILLIAM ACTON, M.R.C.S.** Sixth Edition, 8vo, 12s. [1875]

**URINARY AND REPRODUCTIVE ORGANS :**

their Functional Diseases, by **D. CAMPBELL BLACK, M.D., L.R.C.S.** Edin. Second Edition. 8vo, 10s. 6d. [1875]

**PRACTICAL PATHOLOGY :**

containing Lectures on Suppurative Fever, Diseases of the Veins, Hæmorrhoidal Tumours, Diseases of the Rectum, Syphilis, Gonorrhæal Ophthalmia, &c., by **HENRY LEE, F.R.C.S.**, Surgeon to St. George's Hospital. Third Edition, in 2 vols. 8vo, 10s. each. [1870]

BY THE SAME AUTHOR,

**LECTURES ON SYPHILIS,**

and on some forms of Local Disease, affecting principally the Organs of Generation. With Engravings, 8vo, 10s. [1875]

**SYPHILITIC NERVOUS AFFECTIONS :**

Their Clinical Aspects, by **THOMAS BUZZARD, M.D., F.R.C.P.** Lond., Physician to the National Hospital for Paralysis and Epilepsy. Post 8vo, 5s. [1874]

**PATHOLOGY OF THE URINE,**

including a Complete Guide to its Analysis, by J. L. W. THUDICHUM, M.D., F.R.C.P. Second Edition, rewritten and enlarged, with Engravings, 8vo, 15s. [1877]

**GENITO-URINARY ORGANS, INCLUDING SYPHILIS:**

A Practical Treatise on their Surgical Diseases, designed as a Manual for Students and Practitioners, by W. H. VAN BUREN, M.D., Professor of the Principles of Surgery in Bellevue Hospital Medical College, New York, and E. L. KEYES, M.D., Professor of Dermatology in Bellevue Hospital Medical College, New York. Royal 8vo, with 140 Engravings, 21s. [1874]

**SYPHILIS:**

A Treatise by WALTER J. COULSON, F.R.C.S., Surgeon to the Lock Hospital. 8vo, 10s. [1869]

BY THE SAME AUTHOR,

**STONE IN THE BLADDER:**

Its Prevention, Early Symptoms, and Treatment by Lithotrity. 8vo, 6s. [1868]

**HISTOLOGY AND HISTO-CHEMISTRY OF MAN:**

A Treatise on the Elements of Composition and Structure of the Human Body, by HEINRICH FREY, Professor of Medicine in Zurich. Translated from the Fourth German Edition by ARTHUR E. J. BARKER, Assistant-Surgeon to University College Hospital. And Revised by the Author. 8vo, with 608 Engravings, 21s. [1874]

**HUMAN PHYSIOLOGY:**

A Treatise designed for the Use of Students and Practitioners of Medicine, by JOHN C. DALTON, M.D., Professor of Physiology and Hygiene in the College of Physicians and Surgeons, New York. Sixth Edition, royal 8vo, with 316 Engravings, 20s. [1875]

**HANDBOOK FOR THE PHYSIOLOGICAL LABORATORY,**

by E. KLEIN, M.D., F.R.S., Assistant Professor in the Pathological Laboratory of the Brown Institution, London; J. BURDON-SANDERSON, M.D., F.R.S., Professor of Practical Physiology in University College, London; MICHAEL FOSTER, M.D., F.R.S., Prælector of Physiology in Trinity College, Cambridge; and T. LAUDER BRUNTON, M.D., F.R.S., Lecturer on Materia Medica at St. Bartholomew's Hospital; edited by J. BURDON-SANDERSON. 8vo, with 123 Plates, 24s. [1873]

**PRACTICAL HISTOLOGY:**

By WILLIAM RUTHERFORD, M.D., Professor of the Institutes of Medicine in the University of Edinburgh. Second Edition, with 63 Engravings. Crown 8vo (with additional leaves for notes), 6s. [1876]

**THE MARRIAGE OF NEAR KIN,**

Considered with respect to the Laws of Nations, Results of Experience, and the Teachings of Biology, by ALFRED H. HUTH. 8vo, 14s. [1875]

**PRINCIPLES OF HUMAN PHYSIOLOGY,**

by W. B. CARPENTER, C.B., M.D., F.R.S. Eighth Edition by HENRY POWER, M.B., F.R.C.S., Examiner in Natural Science, University of Oxford, and in Natural Science and Medicine, University of Cambridge. 8vo, with 3 Steel Plates and 371 Engravings, 31s. 6d. [1876]

**STUDENTS' GUIDE TO HUMAN OSTEOLOGY,**

By WILLIAM WARWICK WAGSTAFFE, F.R.C.S., Assistant-Surgeon and Lecturer on Anatomy, St. Thomas's Hospital. With 23 Plates and 66 Engravings. Fcap. 8vo, 10s. 6d. [1875]

**LANDMARKS, MEDICAL AND SURGICAL,**

By LUTHER HOLDEN, F.R.C.S., Senior Surgeon to St. Bartholomew's Hospital. Second Edition, 8vo, 3s. 6d. [1877]

**PATHOLOGICAL ANATOMY :**

Lectures by SAMUEL WILKS, M.D., F.R.S., Physician to, and Lecturer on Medicine at, Guy's Hospital; and WALTER MOXON, M.D., F.R.C.P., Physician to, and Lecturer on Materia Medica at, Guy's Hospital. Second Edition, 8vo, with Plates, 18s. [1875]

**PATHOLOGICAL ANATOMY :**

A Manual by C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital, and EDWARD H. SIEVEKING, M.D., F.R.C.P., Physician to St. Mary's Hospital. Edited by J. F. PAYNE, M.D., F.R.C.P., Assistant Physician and Lecturer on General Pathology at St. Thomas's Hospital. Second Edition, crown 8vo, with 195 Engravings, 16s. [1875]

**POST-MORTEM EXAMINATIONS :**

a Description and Explanation of the Method of Performing them, with especial Reference to Medico-Legal Practice. By Professor RUDOLPH VIRCHOW, of Berlin. Fcap 8vo, 2s. 6d. [1876]

**STUDENT'S GUIDE TO SURGICAL ANATOMY :**

a Text-book for the Pass Examination, by E. BELLAMY, F.R.C.S., Surgeon and Lecturer on Anatomy at Charing Cross Hospital. Fcap 8vo, with 50 Engravings, 6s. 6d. [1873]

**DIAGRAMS OF THE NERVES OF THE HUMAN BODY,**

Exhibiting their Origin, Divisions, and Connexions, with their Distribution, by WILLIAM H. FLOWER, F.R.S., Conservator of Museum, Royal College of Surgeons. Second Edition, roy. 4to, 12s. [1873]

**PRACTICAL ANATOMY :**

a Manual of Dissections by CHRISTOPHER HEATH, F.R.C.S., Surgeon to University College Hospital, and Holme Professor of Surgery in University College. Fourth Edition, crown 8vo, with 16 Coloured Plates and 264 Engravings, 14s. [1877]

**MEDICAL ANATOMY,**

by FRANCIS SIBSON, M.D., F.R.C.P., F.R.S. Imp. folio, with 21 coloured Plates, cloth, 42s., half-morocco, 50s. [1869]

**AN ATLAS OF HUMAN ANATOMY :**

illustrating most of the ordinary Dissections, and many not usually practised by the Student. To be completed in 12 or 13 Bi-monthly Parts, each containing 4 Coloured Plates, with Explanatory Text. By RICKMAN J. GODLEE, M.S., F.R.C.S., Assistant Surgeon to University College Hospital, and Senior Demonstrator of Anatomy in University College. Imp. 4to, 7s. 6d. each Part. [1877-8]

**THE ANATOMIST'S VADE-MECUM :**

a System of Human Anatomy by ERASMUS WILSON, F.R.C.S., F.R.S. Ninth Edition, by G. BUCHANAN, M.A., M.D., Professor of Clinical Surgery in the University of Glasgow, and HENRY E. CLARK, F.F.P.S., Lecturer on Anatomy at the Glasgow Royal Infirmary School of Medicine. Crown 8vo, with 371 Engravings, 14s. [1873]

**ATLAS OF TOPOGRAPHICAL ANATOMY,**

after Plane Sections of Frozen Bodies. By WILHELM BRAUNE, Professor of Anatomy in the University of Leipzig. Translated by EDWARD BELLAMY, F.R.C.S., Surgeon to, and Lecturer on Anatomy, &c., at, Charing Cross Hospital. With 34 Photo-lithographic Plates and 46 Woodcuts. Large Imp. 8vo, 40s. [1877]

**THE STUDENT'S GUIDE TO THE PRACTICE OF MEDICINE,**

by MATTHEW CHARTERIS, M.D., Professor of Medicine in Anderson's College, and Lecturer on Clinical Medicine in the Royal Infirmary, Glasgow. With Engravings on Copper and Wood, fcap. 8vo, 6s. 6d. [1877]

**THE MICROSCOPE IN MEDICINE,**

by LIONEL S. BEALE, M.B., F.R.S., Physician to King's College Hospital. Fourth Edition, with 86 Plates, 8vo, 21s. [1877]

**THE STUDENT'S GUIDE TO MEDICAL DIAGNOSIS,**

by SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Fourth Edition, fcap. 8vo, with 106 Engravings, 6s. 6d. [1876]

**A MANUAL OF MEDICAL DIAGNOSIS,**

by A. W. BARCLAY, M.D., F.R.C.P., Physician to, and Lecturer on Medicine at, St. George's Hospital. Third Edition, fcap 8vo, 10s. 6d. [1876]

**THE ANATOMICAL REMEMBRANCER;**

or, Complete Pocket Anatomist. Eighth Edition, 32mo, 3s. 6d. [1876]

**HOOPER'S PHYSICIAN'S VADE-MECUM ;**

or, Manual of the Principles and Practice of Physic, Ninth Edition by W. A. GUY, M.B., F.R.S., and JOHN HARLEY, M.D., F.R.C.P. Fcap 8vo, with Engravings, 12s. 6d. [1874]

**A NEW SYSTEM OF MEDICINE;**

entitled Recognisant Medicine, or the State of the Sick, by BHOLANOTH BOSE, M.D., Indian Medical Service. 8vo, 10s. 6d. [1877]

BY THE SAME AUTHOR.

**PRINCIPLES OF RATIONAL THERAPEUTICS.**

Commenced as an Inquiry into the Relative Value of Quinine and Arsenic in Ague. 8vo, 4s. 6d. [1877]

**THE MEDICAL REMEMBRANCER;**

or, Book of Emergencies. By E. SHAW, M.R.C.S. Fifth Edition by JONATHAN HUTCHINSON, F.R.C.S., Senior Surgeon to the London Hospital. 32mo, 2s. 6d. [1867]

**MATERIA MEDICA AND THERAPEUTICS:**

(Vegetable Kingdom), by CHARLES D. F. PHILLIPS, M.D., F.R.C.S.E. 8vo, 15s. [1874]

**CLINICAL MEDICINE:**

Lectures and Essays by BALTHAZAR FOSTER, M.D., F.R.C.P. Lond., Professor of Medicine in Queen's College, Birmingham. 8vo, 10s. 6d. [1874]

**CLINICAL STUDIES:**

Illustrated by Cases observed in Hospital and Private Practice, by Sir J. ROSE CORMACK, M.D., F.R.S.E., Physician to the Hertford Hospital of Paris. 2 vols., post 8vo, 20s. [1876]

**ROYLE'S MANUAL OF MATERIA MEDICA AND THERAPEUTICS.**

Sixth Edition by JOHN HARLEY, M.D., F.R.C.P., Assistant Physician to, and Joint Lecturer on Physiology at, St. Thomas's Hospital. Crown 8vo, with 139 Engravings, 15s. [1876]

**PRACTICAL THERAPEUTICS:**

A Manual by E. J. WARING, M.D., F.R.C.P. Lond. Third Edition, fcap 8vo, 12s. 6d. [1871]

**THE ELEMENTS OF THERAPEUTICS.**

A Clinical Guide to the Action of Drugs, by C. BINZ, M.D., Professor of Pharmacology in the University of Bonn. Translated and Edited with Additions, in Conformity with the British and American Pharmacopœias, by EDWARD I. SPARKS, M.A., M.B. Oxon., formerly Radcliffe Travelling Fellow. Crown 8vo, 8s. 6d. [1877]

**THE STUDENT'S GUIDE TO MATERIA MEDICA,**

by JOHN C. THOROWGOOD, M.D., F.R.C.P. Lond., Physician to the City of London Hospital for Diseases of the Chest. Fcap 8vo, with Engravings, 6s. 6d. [1874]

**THE DISEASES OF CHILDREN:**

A Practical Manual, with a Formulary, by EDWARD ELLIS, M.D., late Senior Physician to the Victoria Hospital for Children. Third Edition, crown 8vo, 7s. 6d. [1878]

**THE WASTING DISEASES OF CHILDREN,**

by EUSTACE SMITH, M.D., F.R.C.P. Lond., Physician to the King of the Belgians, Physician to the East London Hospital for Children. Second Edition, post 8vo, 7s. 6d. [1870]

BY THE SAME AUTHOR,

**CLINICAL STUDIES OF DISEASE IN CHILDREN.**

Post 8vo, 7s. 6d. [1876]

**INFANT FEEDING AND ITS INFLUENCE ON LIFE;**

or, the Causes and Prevention of Infant Mortality, by CHARLES H. F. ROUTH, M.D., Senior Physician to the Samaritan Hospital for Women and Children. Third Edition, fcap 8vo, 7s. 6d. [1876]

**COMPENDIUM OF CHILDREN'S DISEASES:**

A Handbook for Practitioners and Students, by JOHANN STEINER, M.D., Professor in the University of Prague. Translated from the Second German Edition by LAWSON TAIT, F.R.C.S., Surgeon to the Birmingham Hospital for Women. 8vo, 12s. 6d. [1874]

**THE DISEASES OF CHILDREN:**

Essays by WILLIAM HENRY DAY, M.D., Physician to the Samaritan Hospital for Diseases of Women and Children. Fcap 8vo, 5s. [1873]

**PUERPERAL DISEASES:**

Clinical Lectures by FORDYCE BARKER, M.D., Obstetric Physician to Bellevue Hospital, New York. 8vo, 15s. [1874]

**THE STUDENT'S GUIDE TO THE PRACTICE OF MIDWIFERY,**

by D. LLOYD ROBERTS, M.D., F.R.C.P., Physician to St. Mary's Hospital, Manchester. Fcap. 8vo, with 95 Engravings, 6s. 6d. [1875]

**OBSTETRIC MEDICINE AND SURGERY,**

Their Principles and Practice, by F. H. RAMSBOTHAM, M.D., F.R.C.P. Fifth Edition, 8vo, with 120 Plates, 22s. [1867]

**OBSTETRIC SURGERY:**

A Complete Handbook, giving Short Rules of Practice in every Emergency, from the Simplest to the most Formidable Operations connected with the Science of Obstetrics, by CHARLES CLAY, Ext.L.R.C.P. Lond., L.R.C.S.E., late Senior Surgeon and Lecturer on Midwifery, St. Mary's Hospital, Manchester. Fcap 8vo, with 91 Engravings, 6s. 6d. [1874]

**SCHROEDER'S MANUAL OF MIDWIFERY,**

including the Pathology of Pregnancy and the Puerperal State. Translated by CHARLES H. CARTER, B.A., M.D. 8vo, with Engravings, 12s. 6d. [1873]

**A HANDBOOK OF UTERINE THERAPEUTICS,**

and of Diseases of Women, by E. J. TILT, M.D., M.R.C.P. Fourth Edition, post-8vo, 10s. [1878]

BY THE SAME AUTHOR,

**THE CHANGE OF LIFE**

in Health and Disease: a Practical Treatise on the Nervous and other Affections incidental to Women at the Decline of Life. Third Edition, 8vo, 10s. 6d. [1870]

**DISEASES OF THE OVARIES:**

their Diagnosis and Treatment, by T. SPENCER WELLS, F.R.C.S., Surgeon to the Queen's Household and to the Samaritan Hospital. 8vo, with about 150 Engravings, 21s. [1873]

**PRACTICAL GYNÆCOLOGY:**

A Handbook of the Diseases of Women, by HEYWOOD SMITH, M.D. Oxon., Physician to the Hospital for Women and to the British Lying-in Hospital. With Engravings, crown 8vo, 5s. 6d. [1877]



**OBSTETRIC OPERATIONS.**

*relating the Treatment of Haemorrhage, and forming a Guide to the Management of Difficult Labour; Lectures by ROBERT BARNES, M.D., F.R.C.P., Consulting Physician and Lecturer on Obstetrics and the Diseases of Women and Children at St. George's Hospital. Third Edition, 1877, with 124 Engravings. 18s.* [1878]

BY THE SAME AUTHOR.

**MEDICAL AND SURGICAL DISEASES OF WOMEN:**

*a Clinical History. Second Edition, 8vo, with 181 Engravings.* [1878]

**OBSTETRIC APHORISMS:**

*for the Use of Students commencing Midwifery Practice by J. G. SWAYNE, M.D., Consulting Physician-Accoucheur to the Bristol General Hospital and Lecturer on Obstetric Medicine at the Bristol Medical School. Sixth Edition, fcap 8vo, with Engravings, 3s. 6d.* [1876]

**RUPTURE OF THE FEMALE PERINEUM,**

*Its treatment, immediate and remote, by GEORGE G. BANTOCK, M.D., Surgeon for In-patients to the Samaritan Free Hospital for Women and Children. With 2 plates, 8vo, 3s. 6d.* [1878]

**HANDBOOK FOR NURSES FOR THE SICK,**

*by ZEPHERINA P. VEITCH. Second Edition, crown 8vo, 3s. 6d.* [1876]

**A MANUAL FOR HOSPITAL NURSES**

*and others engaged in Attending on the Sick by EDWARD J. DOMVILLE, L.R.C.P., M.R.C.S., Surgeon to the Exeter Lying-in Charity. Third Edition, crown 8vo, 2s. 6d.* [1878]

**THE NURSE'S COMPANION:**

*A Manual of General and Monthly Nursing, by CHARLES J. CULLINGWORTH, Surgeon to St. Mary's Hospital, Manchester. Fcap. 8vo, 2s. 6d.* [1876]

**LECTURES ON NURSING,**

*by WILLIAM ROBERT SMITH, M.B., Honorary Medical Officer, Hospital for Sick Children, Sheffield. Second Edition, with 26 Engravings. Post 8vo, 6s.* [1878]

**HOSPITAL MORTALITY**

*being a Statistical Investigation of the Returns of the Hospitals of Great Britain and Ireland for fifteen years, by LAWSON TAIT, F.R.C.S., F.S.S. 8vo, 8s. 6d.* [1877]

**ENGLISH MIDWIVES:**

*their History and Prospects, by J. H. AVELING, M.D., Physician to the Chelsea Hospital for Women, Examiner of Midwives for the Obstetrical Society of London. Crown 8vo, 5s.* [1878]

**A COMPENDIUM OF DOMESTIC MEDICINE**

*and Companion to the Medicine Chest; intended as a Source of Easy Reference for Clergymen, and for Families residing at a Distance from Professional Assistance, by JOHN SAVORY, M.S.A. Ninth Edition, 12mo, 5s.* [1878]

**THE COTTAGE HOSPITAL:**

Its Origin, Progress, Management, and Work, by HENRY C. BUEDETT, the Seaman's Hospital, Greenwich. With Engravings, crown 8vo, 7s. 6d. [1877]

**WINTER COUGH:**

(Catarrh, Bronchitis, Emphysema, Asthma), Lectures by HORACE DOBELL, M.D., Consulting Physician to the Royal Hospital for Diseases of the Chest. Third Edition, with Coloured Plates, 8vo, 1s. 6d. [1875]

BY THE SAME AUTHOR,

**THE TRUE FIRST STAGE OF CONSUMPTION.**

(Lectures.) Crown 8vo, 3s. 6d. [1867]

**DISEASES OF THE CHEST:**

Contributions to their Clinical History, Pathology, and Treatment, by A. T. H. WATERS, M.D., F.R.C.P., Physician to the Liverpool Royal Infirmary. Second Edition, 8vo, with Plates, 15s. [1873]

**NOTES ON ASTHMA;**

its Forms and Treatment, by JOHN C. THOROWGOOD, M.D. Lond., F.R.C.P., Physician to the Hospital for Diseases of the Chest, Victoria Park. Third Edition, crown 8vo, 4s. 6d. [1878]

**ASTHMA**

Its Pathology and Treatment, by J. B. BERKART, M.D., Assistant Physician to the City of London Hospital for Diseases of the Chest. 8vo, 7s. 6d. [1878]

**PROGNOSIS IN CASES OF VALVULAR DISEASE OF THE Heart,** by THOMAS B. PEACOCK, M.D., F.R.C.P., Honorary Consulting Physician to St. Thomas's Hospital. 8vo, 3s. 6d. [1877]

**DISEASES OF THE HEART:**

Their Pathology, Diagnosis, Prognosis, and Treatment (a Manual), by ROBERT H. SEMPLE, M.D., F.R.C.P., Physician to the Hospital for Diseases of the Throat. 8vo, 8s. 6d. [1876]

**PHTHISIS:**

In a series of Clinical Studies, by AUSTIN FLINT, M.D., Professor of the Principles and Practice of Medicine and of Clinical Medicine in the Bellevue Hospital Medical College. 8vo, 16s. [1875]

BY THE SAME AUTHOR,

**A MANUAL OF PERCUSSION AND AUSCULTATION,**

of the Physical Diagnosis of Diseases of the Lungs and Heart, and of Thoracic Aneurism. Post 8vo, 6s. 6d. [1876]

**GROWTHS IN THE LARYNX,**

with Reports and an Analysis of 100 consecutive Cases treated since the Invention of the Laryngoscope by MORELL MACKENZIE, M.D. Lond., M.R.C.P., Physician to the Hospital for Diseases of the Throat. 8vo, with Coloured Plates, 12s. 6d. [1871]

**DISEASES OF THE HEART AND AORTA,**

By THOMAS HAYDEN, F.K.Q.C.P. Irel., Physician to the Mater Misericordiarum Hospital, Dublin. With 80 Engravings. 8vo, 25s. [1875]

**DISEASES OF THE HEART**

and of the Lungs in Connexion therewith—Notes and Observations by THOMAS SHAPTEY, M.D., F.R.C.P. Lond., Senior Physician to the Devon and Exeter Hospital. 8vo, 7s. 6d. [1874]

**DISEASES OF THE HEART AND AORTA:**

Clinical Lectures by GEORGE W. BALFOUR, M.D., F.R.C.P., Physician to, and Lecturer on Clinical Medicine in, the Royal Infirmary, Edinburgh. 8vo, with Engravings, 12s. 6d. [1876]

**PHYSICAL DIAGNOSIS OF DISEASES OF THE HEART.**

Lectures by ARTHUR E. SANSOM, M.D., F.R.C.P., Assistant Physician to the London Hospital. Second Edition, with Engravings, fcap. 8vo, 4s. 6d. [1876]

**TRACHEOTOMY,**

especially in Relation to Diseases of the Larynx and Trachea, by PUGIN THORNTON, M.R.C.S., late Surgeon to the Hospital for Diseases of the Throat. With Photographic Plates and Woodcuts, 8vo, 5s. 6d. [1876]

**SORE THROAT:**

Its Nature, Varieties, and Treatment, including the Connexion between Affections of the Throat and other Diseases. By PROSSER JAMES, M.D., Lecturer on Materia Medica and Therapeutics at the London Hospital, Physician to the Hospital for Diseases of the Throat. Third Edition, with Coloured Plates, 5s. 6d. [1878.]

**SKETCH OF CANNES AND ITS CLIMATE,**

by TH. DE VALCOURT, M.D. Paris, Physician at Cannes. Second Edition, with Photographic View and 6 Meteorological Charts. Crown 8vo, 2s. 6d. [1873]

**WINTER AND SPRING**

on the Shores of the Mediterranean; or, the Genoese Rivas, Italy, Spain, Greece, the Archipelago, Constantinople, Corsica, Sardinia, Sicily, Corfu, Malta, Tunis, Algeria, Smyrna, Asia Minor, with Biarritz and Arcachon, as Winter Climates. By HENRY BENNET, M.D. Fifth Edition, post 8vo, with numerous Plates, Maps, and Engravings, 12s. 6d. [1874]

BY THE SAME AUTHOR,

**TREATMENT OF PULMONARY CONSUMPTION**

by Hygiene, Climate, and Medicine. Second Edition, 8vo, 5s. [1871]

**THE BATH THERMAL WATERS:**

Historical, Social, and Medical, by JOHN KENT SPENDER, M.D., Surgeon to the Mineral Water Hospital, Bath. With an Appendix on the Climate of Bath by the Rev. L. BLOMEFIELD, M.A., F.L.S., F.G.S. 8vo, 7s. 6d. [1877]

**PRINCIPAL HEALTH RESORTS**

of Europe and Africa, and their Use in the Treatment of Chronic Diseases. A Handbook by THOMAS MORE MADDEN, M.D., M.R.I.A., Vice-President of the Dublin Obstetrical Society. 8vo, 10s. [1876]

**ENDEMIC DISEASES OF TROPICAL CLIMATES,**

with their Treatment, by JOHN SULLIVAN, M.D., M.R.C.P. Post 8vo, 6s. [1877]

**FAMILY MEDICINE FOR INDIA :**

A Manual, by WILLIAM J. MOORE, M.D., Surgeon-Major H.M. Indian Medical Service. Published under the Authority of the Government of India. Third Edition, post 8vo, with 60 Engravings. [In the press]

**DISEASES OF TROPICAL CLIMATES**

and their Treatment: with Hints for the Preservation of Health in the Tropics, by JAMES A. HORTON, M.D., Surgeon-Major, Army Medical Department. Post 8vo, 12s. 6d. [1874]

**HEALTH IN INDIA FOR BRITISH WOMEN**

and on the Prevention of Disease in Tropical Climates by EDWARD J. TILT, M.D., Consulting Physician-Accoucheur to the Farringdon General Dispensary. Fourth Edition, crown 8vo, 5s. [1875]

**BURDWAN FEVER,**

or the Epidemic Fever of Lower Bengal (Causes, Symptoms, and Treatment), by GOPAUL CHUNDER ROY, M.D., Surgeon Bengal Establishment. New Edition, 8vo, 5s. [1876]

**BAZAAR MEDICINES OF INDIA**

and Common Medical Plants: Remarks on their Uses, with Full Index of Diseases, indicating their Treatment by these and other Agents procurable throughout India, &c., by EDWARD J. WARING, M.D., F.R.C.P. Lond., Retired Surgeon H.M. Indian Army. Third Edition. Fcap 8vo, 5s. [1875]

**SOME AFFECTIONS OF THE LIVER**

and Intestinal Canal; with Remarks on Ague and its Sequelæ, Scurvy, Purpura, &c., by STEPHEN H. WARD, M.D. Lond., F.R.C.P., Physician to the Seamen's Hospital, Greenwich. 8vo, 7s. [1873]

**DISEASES OF THE LIVER:**

Lettsomian Lectures for 1872 by S. O. HABERSHON, M.D., F.R.C.P., Senior Physician to Guy's Hospital. Post 8vo, 3s. 6d. [1872]

BY THE SAME AUTHOR,

**DISEASES OF THE STOMACH: DYSPEPSIA.**

Second Edition, crown 8vo, 5s.

BY THE SAME AUTHOR,

**PATHOLOGY OF THE PNEUMOGASTRIC NERVE,**

being the Lumleian Lectures for 1876. Post 8vo, 3s. 6d. [1877]

**FUNCTIONAL NERVOUS DISORDERS :**

Studies by C. HANDFIELD JONES, M.B., F.R.C.P., F.R.S., Physician to St. Mary's Hospital. Second Edition, 8vo, 18s. [1876]

**LECTURES ON DISEASES OF THE NERVOUS SYSTEM,**

by SAMUEL WILKS, M.D., F.R.S., Physician to, and Lecturer on Medicine at, Guy's Hospital. 8vo, 15s. [1878]

**NUTRITION IN HEALTH AND DISEASE :**

A Contribution to Hygiene and to Clinical Medicine. By HENRY BENNET, M.D. Third Edition. 8vo, 7s. Cheap Edition, Fcap. 8vo, 2s. 6d. [1877]

**THE STOMACH AND DUODENUM :**

Their Morbid States and their Relations to the Diseases of other Organs, by SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. 8vo, with 10 Plates, 12s. [1868]

**FOOD AND DIETETICS,**

Physiologically and Therapeutically Considered. By FREDERICK W. PAVY, M.D., F.R.S., Physician to Guy's Hospital. Second Edition, 8vo, 15s. [1876]

**HEADACHES :**

their Causes, Nature, and Treatment. By WILLIAM H. DAY, M.D., Physician to the Samaritan Free Hospital for Women and Children. Second Edition, crown 8vo, with Engravings. 6s. 6d. [1878]

**IMPERFECT DIGESTION :**

its Causes and Treatment by ARTHUR LEARED, M.D., F.R.C.P., Senior Physician to the Great Northern Hospital. Sixth Edition, fcap 8vo, 4s. 6d. [1876]

**MEGRIM, SICK-HEADACHE,**

and some Allied Disorders : a Contribution to the Pathology of Nerve-Storms, by EDWARD LIVEING, M.D. Cantab., F.R.C.P., Hon. Fellow of King's College, London. 8vo, with Coloured Plate, 15s. [1873]

**NEURALGIA AND KINDRED DISEASES**

of the Nervous System : their Nature, Causes, and Treatment, with a series of Cases, by JOHN CHAPMAN, M.D., M.R.C.P. 8vo, 14s. [1873]

**THE SYMPATHETIC SYSTEM OF NERVES,**

and their Functions as a Physiological Basis for a Rational System of Therapeutics by EDWARD MERYON, M.D., F.R.C.P., Physician to the Hospital for Diseases of the Nervous System. 8vo, 3s. 6d. [1873]

**RHEUMATIC GOUT,**

or Chronic Rheumatic Arthritis of all the Joints; a Treatise by ROBERT ADAMS, M.D., M.R.I.A., late Surgeon to H.M. the Queen in Ireland, and Regius Professor of Surgery in the University of Dublin. Second Edition, 8vo, with Atlas of Plates, 21s. [1873]

**GOUT, RHEUMATISM,**

and the Allied Affections; a Treatise by PETER HOOD, M.D. Crown 8vo, 10s. 6d. [1871]

**RHEUMATISM :**

Notes by JULIUS POLLOCK, M.D., F.R.C.P., Senior Physician to, and Lecturer on Medicine at, Charing Cross Hospital. Fcap. 8vo, 2s. 6d. [1878.]

**CANCER :**

its varieties, their Histology and Diagnosis, by HENRY ARNOTT, F.R.C.S., late Assistant-Surgeon to, and Lecturer on Morbid Anatomy at, St. Thomas's Hospital. 8vo, with 5 Plates and 22 Engravings, 5s. 6d. [1872.]

**CANCEROUS AND OTHER INTRA-THORACIC GROWTHS :**

their Natural History and Diagnosis, by J. RISDON BENNETT, M.D., F.R.C.P., Member of the General Medical Council. Post 8vo, with Plates, 8s. [1872.]

**CERTAIN FORMS OF CANCER,**

with a New and successful Mode of Treating it, to which is prefixed a Practical and Systematic Description of all the varieties of this Disease, by ALEX. MARSDEN, M.D., F.R.C.S.E., Consulting Surgeon to the Royal Free Hospital, and Senior Surgeon to the Cancer Hospital. Second Edition, with Coloured Plates, 8vo, 8s. 6d. [1873.]

**ATLAS OF SKIN DISEASES :**

a series of Illustrations, with Descriptive Text and Notes upon Treatment. By TILBURY FOX, M.D., F.R.C.P., Physician to the Department for Skin Diseases in University College Hospital. With 72 Coloured Plates, royal 4to, half morocco, £6 6s. [1877.]

**DISEASES OF THE SKIN :**

a System of Cutaneous Medicine by ERASMUS WILSON, F.R.C.S., F.R.S. Sixth Edition, 8vo, 18s., with Coloured Plates, 36s. [1867.]

BY THE SAME AUTHOR,

**LECTURES ON EKZEMA**

and Ekzematous Affections: with an Introduction on the General Pathology of the Skin, and an Appendix of Essays and Cases. 8vo, 10s. 6d. [1870.]

ALSO,

**LECTURES ON DERMATOLOGY :**

delivered at the Royal College of Surgeons, 1870, 6s. ; 1871-3, 10s. 6d., 1874-5, 10s. 6d.

**ECZEMA :**

by MCCALL ANDERSON, M.D., Professor of Clinical Medicine in the University of Glasgow. Third Edition, 8vo, with Engravings, 7s. 6d. [1874.]

BY THE SAME AUTHOR,

**PARASITIC AFFECTIONS OF THE SKIN**

Second Edition, 8vo, with Engravings, 7s. 6d. [1868.]

**PSORIASIS OR LEPRO,**

by GEORGE GASKOIN, M.R.C.S., Surgeon to the British Hospital for Diseases of the Skin. 8vo, 5s. [1875.]

**MYCETOMA ;**

or, the Fungus Disease of India, by H. VANDYKE CARTER, M.D., Surgeon-Major H.M. Indian Army. 4to, with 11 Coloured Plates, 42s. [1874]

**CERTAIN ENDEMIC SKIN AND OTHER DISEASES**

of India and Hot Climates generally, by TILBURY FOX, M.D., F.R.C.P., and T. FARQUHAR, M.D. (Published under the sanction of the Secretary of State for India in Council). 8vo, 10s. 6d. [1876]

**DISEASES OF THE SKIN,**

in Twenty-four Letters on the Principles and Practice of Cutaneous Medicine, by HENRY EVANS CAUTY, M.R.C.S., Surgeon to the Liverpool Dispensary for Diseases of the Skin, 8vo, 12s. 6d. [1874]

**THE HAIR IN HEALTH AND DISEASE,**

by E. WYNDHAM COTTLE, F.R.C.S., Senior Assistant Surgeon to the Hospital for Diseases of the Skin, Blackfriars. Fcap. 8vo, 2s. 6d. [1877]

**WORMS:**

a Series of Lectures delivered at the Middlesex Hospital on Practical Helminthology by T. SPENCER COBBOLD, M.D., F.R.S. Post 8vo, 5s. [1872]

**THE LAWS AFFECTING MEDICAL MEN :**

a Manual by ROBERT G. GLENN, LL.B., Barrister-at-Law; with a Chapter on Medical Etiquette by Dr. A. CARPENTER. 8vo, 14s. [1871]

**MEDICAL JURISPRUDENCE,**

Its Principles and Practice, by ALFRED S. TAYLOR, M.D., F.R.C.P., F.R.S. Second Edition, 2 vols., 8vo, with 189 Engravings, £1 11s. 6d. [1873]

BY THE SAME AUTHOR,

**A MANUAL OF MEDICAL JURISPRUDENCE.**

Ninth Edition. Crown 8vo, with Engravings, 14s. [1874]

ALSO,

**POISONS,**

in Relation to Medical Jurisprudence and Medicine. Third Edition, crown 8vo, with 104 Engravings, 16s. [1876]

**MEDICAL JURISPRUDENCE :**

Lectures by FRANCIS OGSTON, M.D., Professor of Medical Jurisprudence and Medical Logic in the University of Aberdeen. Edited by FRANCIS OGSTON, Jun., M.D., Assistant to the Professor of Medical Jurisprudence and Lecturer on Practical Toxicology in the University of Aberdeen. 8vo, with 12 Copper Plates, 18s. [1878]

**A TOXICOLOGICAL CHART,**

exhibiting at one View the Symptoms, Treatment, and mode of Detecting the various Poisons—Mineral, Vegetable, and Animal: with Concise Directions for the Treatment of Suspended Animation, by WILLIAM STOWE, M.R.C.S.E. Thirteenth Edition, 2s.; on roller, 5s. [1872]

**A HANDY-BOOK OF FORENSIC MEDICINE AND TOXICOLOGY,**  
by W. BATHURST WOODMAN, M.D., F.R.C.P., Assistant Physician  
and Co-Lecturer on Physiology and Histology at the London Hospital;  
and C. MEYMOTT TIDY, M.D., F.C.S., Professor of Chemistry and of  
Medical Jurisprudence and Public Health at the London Hospital.  
With 8 Lithographic Plates and 116 Engravings, 8vo, 31s. 6d. [1877]

**THE MEDICAL ADVISER IN LIFE ASSURANCE,**  
by EDWARD HENRY SIEVEKING, M.D., F.R.C.P., Physician to St.  
Mary's and the Lock Hospitals; Physician-Extraordinary to the  
Queen; Physician-in-Ordinary to the Prince of Wales, &c. Crown  
8vo, 6s. [1874]

**IDIOCY AND IMBECILITY,**  
by WILLIAM W. IRELAND, M.D., Medical Superintendent of the  
Scottish National Institution for the Education of Imbecile Children  
at Larbert, Stirlingshire. With Engravings, 8vo, 14s. [1877]

**PSYCHOLOGICAL MEDICINE:**  
a Manual, containing the Lunacy Laws, the Nosology, Ætiology,  
Statistics, Description, Diagnosis, Pathology (including Morbid His-  
tology), and Treatment of Insanity, by J. C. BUCKNILL, M.D.,  
F.R.S., and D. H. TUKE, M.D., F.R.C.P. Third Edition, 8vo, with  
10 Plates and 34 Engravings, 25s. [1873]

**MADNESS:**  
in its Medical, Legal, and Social Aspects, Lectures by EDGAR  
SHEPPARD, M.D., M.R.C.P., Professor of Psychological Medicine in  
King's College; one of the Medical Superintendents of the Colney  
Hatch Lunatic Asylum. 8vo, 6s. 6d. [1873]

**HANDBOOK OF LAW AND LUNACY;**  
or, the Medical Practitioner's Complete Guide in all Matters relating  
to Lunacy Practice, by J. T. SABBEN, M.D., and J. H. BALFOUR  
BROWNE, Barrister-at-Law. 8vo, 5s. [1872]

**INFLUENCE OF THE MIND UPON THE BODY**  
in Health and Disease, Illustrations designed to elucidate the Action  
of the Imagination, by DANIEL HACK TUKE, M.D., F.R.C.P.  
8vo, 14s. [1872]

**A MANUAL OF PRACTICAL HYGIENE,**  
by E. A. PARKES, M.D., F.R.S. Fifth Edition, by F. DE CHAUMONT,  
M.D., Professor of Military Hygiene in the Army Medical School.  
8vo, with 9 Plates and 112 Engravings, 18s. [1878]

**A HANDBOOK OF HYGIENE AND SANITARY SCIENCE,**  
by GEORGE WILSON, M.A., M.D., Medical Officer of Health for Mid-  
Warwickshire. Third Edition, post 8vo, with Engravings, 10s. 6d.  
[1877]

**MICROSCOPICAL EXAMINATION OF DRINKING WATER:**  
A Guide, by JOHN D. MACDONALD, M.D., F.R.S., Assistant Pro-  
fessor of Naval Hygiene, Army Medical School. 8vo, with 24 Plates,  
7s. 6d. [1875]



**HANDBOOK OF MEDICAL AND SURGICAL ELECTRICITY,**

by HERBERT TIBBITS, M.D., F.R.C.P.E., Medical Superintendent of the National Hospital for the Paralysed and Epileptic. Second Edition 8vo, with 95 Engravings, 9s. [1877]

BY THE SAME AUTHOR.

**A MAP OF ZIEMSSSEN'S MOTOR POINTS OF THE HUMAN BODY:**

a Guide to Localised Electrification. Mounted on Rollers, 35 × 21. With 20 Illustrations, 5s. [1877]

**CLINICAL USES OF ELECTRICITY;**

Lectures delivered at University College Hospital by J. RUSSELL REYNOLDS, M.D. Lond., F.R.C.P., F.R.S., Professor of Medicine in University College. Second Edition, post 8vo, 3s. 6d. [1878]

**MEDICO-ELECTRIC APPARATUS:**

A Practical Description of every Form in Modern Use, with Plain Directions for Mounting, Charging, and Working, by SALT & SON, Birmingham. Second Edition, revised and enlarged, with 33 Engravings, 8vo, 2s. 6d. [1877]

**A DICTIONARY OF MEDICAL SCIENCE;**

containing a concise explanation of the various subjects and terms of Medicine, &c.; Notices of Climate and Mineral Waters; Formulæ for Official, Empirical, and Dietetic Preparations; with the Accentuation and Etymology of the terms and the French and other Synonyms, by ROBLEY DUNGLISON, M.D., LL.D. New Edition, royal 8vo, 28s. [1874]

**A MEDICAL VOCABULARY;**

being an Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, giving their derivation, meaning, application, and pronunciation, by ROBERT G. MAYNE, M.D., LL.D. Fourth Edition, fcap 8vo, 10s. [1875]

**ATLAS OF OPHTHALMOSCOPY,**

by R. LIEBREICH, Ophthalmic Surgeon to St. Thomas's Hospital. Translated into English by H. ROSBOROUGH SWANZY, M.B. Dub. Second Edition, containing 59 Figures, 4to, £1 10s. [1870]

**DISEASES OF THE EYE:**

a Manual by C. MACNAMARA, F.R.C.S., Surgeon to Westminster Hospital. Third Edition, fcap. 8vo, with Coloured Plates and Engravings, 12s. 6d. [1876]

**DISEASES OF THE EYE:**

A Practical Treatise by HAYNES WALTON, F.R.C.S., Surgeon to St. Mary's Hospital and in charge of its Ophthalmological Department. Third Edition, 8vo, with 3 Plates and nearly 300 Engravings, 25s. [1875]

**HINTS ON OPHTHALMIC OUT-PATIENT PRACTICE,**

by CHARLES HIGGINS, F.R.C.S., Ophthalmic Assistant Surgeon to, and Lecturer on Ophthalmology at, Guy's Hospital. 87 pp., fcap. 8vo, 2s. 6d. [1877]

**OPHTHALMIC MEDICINE AND SURGERY:**

a Manual by T. WHARTON JONES, F.R.C.S., F.R.S., Professor of Ophthalmic Medicine and Surgery in University College. Third Edition, fcap. 8vo, with 9 Coloured Plates and 173 Engravings, 12s. 6d. [1865]

**DISEASES OF THE EYE:**

A Treatise by J. SOELBERG WELLS, F.R.C.S., Ophthalmic Surgeon to King's College Hospital and Surgeon to the Royal London Ophthalmic Hospital. Third Edition, 8vo, with Coloured Plates and Engravings, 25s. [1873]

BY THE SAME AUTHOR,

**LONG, SHORT, AND WEAK SIGHT,**

and their Treatment by the Scientific use of Spectacles. Fourth Edition, 8vo, 6s. [1873]

**A SYSTEM OF DENTAL SURGERY,**

by JOHN TOMES, F.R.S., and CHARLES S. TOMES, M.A., F.R.S., Lecturer on Dental Anatomy and Physiology at the Dental Hospital of London. Second Edition, fcap 8vo, with 268 Engravings, 14s. [1873]

**DENTAL ANATOMY, HUMAN AND COMPARATIVE:**

A Manual, by CHARLES S. TOMES, M.A., F.R.S., Lecturer on Dental Anatomy and Physiology at the Dental Hospital of London. With 179 Engravings, crown 8vo, 10s. 6d. [1876]

**A MANUAL OF DENTAL MECHANICS,**

with an Account of the Materials and Appliances used in Mechanical Dentistry, by OAKLEY COLES, L.D.S., R.C.S., Surgeon-Dentist to the Hospital for Diseases of the Throat. Second Edition, crown 8vo, with 140 Engravings, 7s. 6d. [1876]

**HANDBOOK OF DENTAL ANATOMY**

and Surgery for the use of Students and Practitioners by JOHN SMITH, M.D., F.R.S. Edin., Surgeon-Dentist to the Queen in Scotland. Second Edition, fcap 8vo, 4s. 6d. [1871]

**STUDENT'S GUIDE TO DENTAL ANATOMY AND SURGERY,**

by HENRY SEWILL, M.R.C.S., L.D.S., Dentist to the West London Hospital. With 77 Engravings, fcap. 8vo, 5s. 6d. [1876]

**OPERATIVE DENTISTRY:**

A Practical Treatise, by JONATHAN TAFT, D.D.S., Professor of Operative Dentistry in the Ohio College of Dental Surgery. Third Edition, thoroughly revised, with many additions, and 134 Engravings, 8vo, 18s. [1877]

**DENTAL CARIES**

and its Causes: an Investigation into the influence of Fungi in the Destruction of the Teeth, by Drs. LEBER and ROTTENSTEIN. Translated by H. CHANDLER, D.M.D., Professor in the Dental School of Harvard University. With Illustrations, royal 8vo, 5s. [1878.]

**EPIDEMIOLOGY;**

or, the Remote Cause of Epidemic Diseases in the Animal and in the Vegetable Creation, by JOHN PARKIN, M.D., F.R.C.P.E. Part I, Contagion—Modern Theories—Cholera—Epizootics. 8vo, 5s. [1873]

on

The following CATALOGUES issued by Messrs CHURCHILL  
will be forwarded post free on application :

1. *Messrs Churchill's General List of nearly 600 works on Medicine, Surgery, Midwifery, Materia Medica, Hygiene, Anatomy, Physiology, Chemistry, &c., &c., with a complete Index to their Titles, for easy reference.*

N.B.—*This List includes Nos. 2 and 3.*

2. *Selection from Messrs Churchill's General List, comprising all recent Works published by them on the Art and Science of Medicine.*

3. *A descriptive List of Messrs Churchill's Works on Chemistry, Pharmacy, Botany, Photography, Zoology, and other branches of Science.*

4. *Messrs Churchill's Red-Letter List, giving the Titles of forthcoming New Works and New Editions.*

[Published every October.]

5. *The Medical Intelligencer, an Annual List of New Works and New Editions published by Messrs J. & A. Churchill, together with Particulars of the Periodicals issued from their House.*

[Sent in January of each year to every Medical Practitioner in the United Kingdom whose name and address can be ascertained. A large number are also sent to the United States of America, Continental Europe, India, and the Colonies.]

---

MESSRS CHURCHILL have a special arrangement with MESSRS LINDSAY & BLAKISTON, OF PHILADELPHIA, in accordance with which that Firm act as their Agents for the United States of America, either keeping in Stock most of Messrs CHURCHILL's Books, or reprinting them on Terms advantageous to Authors. Many of the Works in this Catalogue may therefore be easily obtained in America.



